



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/54, 15/82, 9/10, A01H 5/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/66747</b>																														
			(43) International Publication Date: 9 November 2000 (09.11.00)																														
(21) International Application Number: PCT/GB00/01572 (22) International Filing Date: 20 April 2000 (20.04.00) (30) Priority Data: <table border="0"> <tr><td>9917835.2</td><td>29 April 1999 (29.04.99)</td><td>GB</td></tr> <tr><td>9909972.3</td><td>29 April 1999 (29.04.99)</td><td>GB</td></tr> <tr><td>9909967.3</td><td>29 April 1999 (29.04.99)</td><td>GB</td></tr> <tr><td>9909981.4</td><td>29 April 1999 (29.04.99)</td><td>GB</td></tr> <tr><td>9909969.9</td><td>29 April 1999 (29.04.99)</td><td>GB</td></tr> <tr><td>9917843.6</td><td>29 July 1999 (29.07.99)</td><td>GB</td></tr> <tr><td>9917836.0</td><td>29 July 1999 (29.07.99)</td><td>GB</td></tr> <tr><td>9930212.7</td><td>21 December 1999 (21.12.99)</td><td>GB</td></tr> <tr><td>9930210.1</td><td>21 December 1999 (21.12.99)</td><td>GB</td></tr> <tr><td>9930202.8</td><td>21 December 1999 (21.12.99)</td><td>GB</td></tr> </table> (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): HAWKES, Timothy, Robert [GB/GB]; Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). WARNER, Simon, Anthony, James [GB/GB]; Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB).		9917835.2	29 April 1999 (29.04.99)	GB	9909972.3	29 April 1999 (29.04.99)	GB	9909967.3	29 April 1999 (29.04.99)	GB	9909981.4	29 April 1999 (29.04.99)	GB	9909969.9	29 April 1999 (29.04.99)	GB	9917843.6	29 July 1999 (29.07.99)	GB	9917836.0	29 July 1999 (29.07.99)	GB	9930212.7	21 December 1999 (21.12.99)	GB	9930210.1	21 December 1999 (21.12.99)	GB	9930202.8	21 December 1999 (21.12.99)	GB	ANDREWS, Christopher, John [GB/GB]; Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). BACHOO, Satvinder [GB/GB]; Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). PICKERILL, Andrew, Paul [GB/GB]; Jealott's Hill International Research Centre, Bracknell, Berkshire RG42 6ET (GB). (74) Agents: GAAL, Jozsef, Christopher et al.; Zeneca Agrochemicals, Intellectual Property Dept., P.O. Box 3538, Jealott's Hill Research Station, Bracknell RG42 6YA (GB). (81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
9917835.2	29 April 1999 (29.04.99)	GB																															
9909972.3	29 April 1999 (29.04.99)	GB																															
9909967.3	29 April 1999 (29.04.99)	GB																															
9909981.4	29 April 1999 (29.04.99)	GB																															
9909969.9	29 April 1999 (29.04.99)	GB																															
9917843.6	29 July 1999 (29.07.99)	GB																															
9917836.0	29 July 1999 (29.07.99)	GB																															
9930212.7	21 December 1999 (21.12.99)	GB																															
9930210.1	21 December 1999 (21.12.99)	GB																															
9930202.8	21 December 1999 (21.12.99)	GB																															
(54) Title: HERBICIDE RESISTANT PLANTS																																	
(57) Abstract																																	
<p>The present invention provides, inter alia, an isolated polynucleotide comprising a region encoding a chloroplast transit peptide and a glyphosate resistant 5-enolpyruvylshikimate phosphate synthase (EPSPS) 3' of the peptide, the said region being under expression control of a plant operable promoter, with the provisos that the said promoter is not heterologous with respect to the said region, and the chloroplast transit peptide is not heterologous with respect to the said synthase.</p>																																	



## HERBICIDE RESISTANT PLANTS

The present invention relates to recombinant DNA technology, and in particular to the production of transgenic plants which exhibit substantial resistance or substantial  
5 tolerance to herbicides when compared with non transgenic like plants. The invention also relates, *inter alia*, to the nucleotide sequences (and expression products thereof) which are used in the production of, or are produced by, the said transgenic plants.

Plants which are substantially "tolerant" to a herbicide when they are subjected to it provide a dose/response curve which is shifted to the right when compared with that  
10 provided by similarly subjected non tolerant like plants. Such dose/response curves have "dose" plotted on the x-axis and "percentage kill", "herbicidal effect" etc. plotted on the y-axis. Tolerant plants will typically require at least twice as much herbicide as non tolerant like plants in order to produce a given herbicidal effect. Plants which are substantially "resistant" to the herbicide exhibit few, if any, necrotic, lytic, chlorotic or other lesions when  
15 subjected to the herbicide at concentrations and rates which are typically employed by the agricultural community to kill weeds in the field in which crops are to be grown for commercial purposes.

It is particularly preferred that the plants are substantially resistant or substantially tolerant to herbicides (hereinafter "glyphosate") which have 5-enol pyruvyl shikimate  
20 phosphate synthetase (hereinafter "EPSPS") as their site of action, of which N-phosphonomethylglycine (and its various salts) is the pre-eminent example.

The herbicide may be applied either pre- or post emergence in accordance with usual techniques for herbicide application to fields comprising crops which have been rendered resistant to the herbicide. The present invention provides, *inter alia*, nucleotide sequences  
25 useful in the production of such herbicide tolerant or resistant plants.

According to the present invention there is provided an isolated polynucleotide comprising the sequence depicted in SEQ ID No.41. The invention also provides a polynucleotide, excluding the cDNA encoding the rice and corn EPSPS, which encodes an EPSPS and which is complementary to one which when incubated at a temperature of  
30 between 65 and 70°C in 0.1 strength citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.1 strength citrate buffered saline containing 0.1% SDS still hybridises with the sequence depicted in SEQ ID No. 41. An EPSPS encoding

polynucleotide according to the invention may, however, be obtained by screening plant genomic DNA libraries with a nucleotide constituting an intron within the SEQ ID No. 41 sequence, and the invention also includes such a sequence obtainable from that screening.

The invention also includes an isolated polynucleotide comprising a region encoding a chloroplast transit peptide and a glyphosate resistant 5-enolpyruvylshikimate phosphate synthase (EPSPS) 3' of the peptide, the said region being under expression control of a plant operable promoter, with the *provisos* that the said promoter is not heterologous with respect to the said region, and the chloroplast transit peptide is not heterologous with respect to the said synthase.

By "heterologous" is meant from a different source, and correspondingly "non-heterologous" means derived from the same source - but at a gene rather than organism or tissue level. For example the CaMV35S promoter is clearly heterologous with respect to a petunia EPSPS coding sequence insofar as the promoter is derived from a virus and the sequence - the expression of which it controls - from a plant. The term "heterologous" according to the present invention has a still narrower meaning, however. For example "heterologous" as it relates to the present invention means that the petunia EPSPS coding sequence is "heterologous" with respect to, for example, a promoter also derived from petunia - other than that which controls expression of the EPSPS gene. In this sense the petunia promoter derived from the petunia EPSPS gene then used to control expression of an EPSPS coding sequence likewise-derived from petunia is "non-heterologous" with respect to the said coding sequence. "Non-heterologous" does not mean, however, that the promoter and coding sequence must necessarily have been obtained from one and the same (original or progenitor) polynucleotide. Likewise with respect to transit peptides. For example, a rubisco chloroplast transit peptide derived from sunflower is "heterologous" with respect to the coding sequence of an EPSPS gene likewise derived from sunflower (the same plant, tissue or cell). A rubisco transit peptide encoding sequence derived from sunflower is "non-heterologous" with respect to a rubisco enzyme encoding-sequence also derived from sunflower even if the origins of both sequences are different polynucleotides which may have been present in different cells, tissues or sunflower plants.

A preferred form of the polynucleotide comprises the following components in the 5' to 3' direction of transcription:-

- (i) At least one transcriptional enhancer being that enhancing region which is upstream from the transcriptional start of the sequence from which the enhancer is obtained and which enhancer *per se* does not function as a promoter either in the sequence in which it is endogenously comprised or when present heterologously as part of a construct;
- (ii) The promoter from the rice EPSPS gene;
- (iii) The rice genomic sequence which encodes the rice EPSPS chloroplast transit peptide;
- (iv) The genomic sequence which encodes the rice EPSPS;
- (v) A transcriptional terminator;
- wherein the rice EPSPS coding sequence is modified in that a first position is mutated so that the residue at this position is Ile rather than Thr and a second position is mutated so that the residue at this position is Ser rather than Pro, the mutations being introduced into EPSPS sequences which comprise the following conserved region GNAGTAMRPLTAAV in the wild type enzyme such that modified sequence reads GNAGIAMRSLTAAV.

The enhancing region preferably comprises a sequence the 3' end of which is at least 40 nucleotides upstream of the closest transcriptional start of the sequence from which the enhancer is obtained. In a further embodiment of the polynucleotide, the enhancing region comprises a region the 3' end of which is at least 60 nucleotides upstream of the said closest start, and in a still further embodiment of the polynucleotide the said enhancing region comprises a sequence the 3' end of which is at least 10 nucleotides upstream from the first nucleotide of the TATA consensus of the sequence from which the enhancer is obtained.

The polynucleotide according to the invention may comprise two or more transcriptional enhancers, which in a particular embodiment of the polynucleotide may be tandemly present.

In the present inventive polynucleotide the 3' end of the enhancer, or first enhancer if there is more than one present, may be between about 100 to about 1000 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide or the first nucleotide of an intron in the 5' untranslated region in the case that the said region contains an intron. In a more preferred embodiment of the polynucleotide, the 3' end of the enhancer, or first enhancer, is between about 150 to about 1000 nucleotides upstream of the

codon corresponding to the translational start of the EPSPS transit peptide or the first nucleotide of an intron in the 5' untranslated region, and in a still more preferred embodiment the 3' end of the enhancer, or first enhancer, may be between about 300 to about 950 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide or the first nucleotide of an intron in the 5' untranslated region. In a yet more preferred embodiment, the 3' end of the enhancer, or first enhancer, may be located between about 770 and about 790 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide or the first nucleotide of an intron in the 5' untranslated region.

10 In an alternative inventive polynucleotide, the 3' end of the enhancer, or first enhancer, may be located between about 300 to about 380 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide or the first nucleotide of an intron in the 5' untranslated region, and in a preferred embodiment of this alternative polynucleotide the 3' end of the enhancer, or first enhancer, is located between about 320 to  
15 about 350 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.

In the polynucleotide according to the invention, the region upstream of the promoter from the rice EPSPS gene may comprise at least one enhancer derived from a sequence which is upstream from the transcriptional start of either the maize polyubiquitin or rice actin  
20 promoters.

Accordingly the polynucleotide may comprise in the 5' to 3' direction a first enhancer comprising a transcriptional enhancing region derived from a sequence which is upstream from the transcriptional start of either the rice actin promoter and a second enhancer comprising a transcriptional enhancing region derived from a sequence which is upstream  
25 from the transcriptional start of the rice actin promoter.

Whatever the identity and juxtaposition of the various enhancers present in the polynucleotide, the nucleotides 5' of the codon which constitutes the translational start of the rice EPSPS chloroplast transit peptide may be Kozack preferred. The skilled man will be aware of what is meant by this - which in any event will be further apparent from the  
30 following examples. Particularly preferred embodiments of the present inventive polynucleotide have a non-translated region which comprises a sequence which functions as an intron located 5' of the rice genomic sequence which encodes the rice EPSPS chloroplast

transit peptide. The non-translated region may comprise the sequence depicted in SEQ ID NO. 48.

The polynucleotide of the invention may comprise a virally derived translational enhancer located within the non translated region 5' of the rice genomic sequence which encodes the rice EPSPS chloroplast transit peptide. The man skilled in the art is aware of the identity of such suitable translational enhancers - such as the Omega and Omega prime sequences derived from TMV and that derived from the tobacco etch virus, and how such translational enhancers can be introduced into the polynucleotide so as to provide for the desired result.

The polynucleotide according to the invention may further comprise regions encoding proteins capable of conferring upon plant material containing it at least one of the following agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, dessication, and herbicides. Whilst such a polynucleotide contemplates the herbicide resistance conferring gene being other than an EPSPS, such as glyphosate oxido-reductase (GOX) for example, the herbicide may be other than glyphosate in which case the resistance conferring genes may be selected from the group encoding the following proteins: phosphinothricin acetyl transferase (PAT), hydroxyphenyl pyruvate dioxygenase (HPPD), glutathione S transferase (GST), cytochrome P450, Acetyl-CoA carboxylase (ACCase), Acetolactate synthase (ALS), protoporphyrinogen oxidase (PPO), dihydropteroate synthase, polyamine transport proteins, superoxide dismutase (SOD), bromoxynil nitrilase, phytoene desaturase (PDS), the product of the *ifdA* gene obtainable from *Alcaligenes eutrophus*, and known mutagenised or otherwise modified variants of the said proteins. In the case that the polynucleotide provides for multiple herbicide resistance such herbicides may be selected from the group consisting of a dinitroaniline herbicide, triazolo-pyrimidines, uracil, a phenylurea, triketone, isoxazole, acetanilide, oxadiazole, triazinone, sulfonanilide, amide, anilide, RP201772, flurochloridone, norflurazon, and triazolinone type herbicide and the post-emergence herbicide is selected from the group consisting of glyphosate and salts thereof, glufosinate, asulam, bentazon, bialaphos, bromacil, sethoxydim or another cyclohexanedione, dicamba, fosamine, flupoxam, phenoxy propionate, quizalofop or another aryloxy-phenoxypropanoate, picloram, fluormetron, atrazine or another triazine, metribuzin, chlorimuron, chlorsulfuron, flumetsulam, halosulfuron, sulfometron, imazaquin, imazethapyr, isoxaben, imazamox, metosulam, pyriethroba, rimsulfuron, bensulfuron, nicosulfuron,

fomesafen, fluroglycofen, KIH9201, ET751, carfentrazone, ZA1296, sulcotrione, paraquat, diquat, bromoxynil and fenoxaprop.

In the case that the polynucleotide comprises sequences encoding insecticidal proteins, these proteins may be selected from the group consisting of crystal toxins derived from Bt, including secreted Bt toxins; protease inhibitors, lectins, Xenhorabodus/Photorhabdus toxins; the fungus resistance conferring genes may be selected from the group consisting of those encoding known AFPs, defensins, chitinases, glucanases, Avr-Cf9. Particularly preferred insecticidal proteins are cryIAC, cryIAB, cry3A, Vip 1A, Vip 1B, cysteine protease inhibitors, and snowdrop lectin. In the case that the polynucleotide comprises bacterial resistance conferring genes these may be selected from the group consisting of those encoding cecropins and techyplesin and analogues thereof. Virus resistance conferring genes may be selected from the group consisting of those encoding virus coat proteins, movement proteins, viral replicases, and anti-sense and ribozyme sequences which are known to provide for virus resistance; whereas the stress, salt, and drought resistance conferring genes may be selected from those that encode Glutathione-S-transferase and peroxidase, the sequence which constitutes the known CBF1 regulatory sequence and genes which are known to provide for accumulation of trehalose.

The polynucleotide according to the invention may be modified to enhance expression of the protein encoding sequences comprised by it, in that mRNA instability motifs and/or fortuitous splice regions may be removed, or crop preferred codons may be used so that expression of the thus modified polynucleotide in a plant yields substantially similar protein having a substantially similar activity/function to that obtained by expression of the unmodified polynucleotide in the organism in which the protein encoding regions of the unmodified polynucleotide are endogenous. The degree of identity between the modified polynucleotide and a polynucleotide endogenously contained within the said plant and encoding substantially the same protein may be such as to prevent co-suppression between the modified and endogenous sequences. In this case the degree of identity between the sequences should preferably be less than about 70%.

The invention still further includes a biological or transformation vector comprising the present inventive polynucleotide. Accordingly, by "vector" is meant, *inter alia*, one of the following: a plasmid, virus, cosmid or a bacterium transformed or transfected so as to contain the polynucleotide.



The invention still further includes plant material which has been transformed with the said polynucleotide or vector, as well as such transformed plant material which has been, or is, further transformed with a polynucleotide comprising regions encoding proteins capable of conferring upon plant material containing it at least one of the following  
5 agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, dessication, and herbicides.

The invention still further includes morphologically normal, fertile whole plants which have been regenerated from the material disclosed in the immediately preceding paragraph, their progeny seeds and parts, which progeny comprises the polynucleotide or  
10 vector of the invention stably incorporated into its genome and heritable in a Mendelian manner.

The invention still further includes morphologically normal fertile whole plants which contain the present inventive polynucleotide and which result from the crossing of plants which have been regenerated from material transformed with the present inventive  
15 polynucleotide or vector, and plants which have been transformed with a polynucleotide comprising regions encoding proteins capable of conferring upon plant material containing it at least one of the following agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, dessication, and herbicides, the progeny of the resultant plants, their seeds and parts.

20 Plants of the invention may be selected from the group consisting of field crops, fruits and vegetables such as canola, sunflower, tobacco, sugar beet, cotton, maize, wheat, barley, rice, sorghum, tomato, mango, peach, apple, pear, strawberry, banana, melon, potato, carrot, lettuce, cabbage, onion, soya spp, sugar cane, pea, field beans, poplar, grape, citrus, alfalfa, rye, oats, turf and forage grasses, flax and oilseed rape, and nut producing plants  
25 insofar as they are not already specifically mentioned, their progeny, seeds and parts.

Particularly preferred such plants include maize, soybean, cotton, sugar beet and canola.

The invention still further comprises a method of selectively controlling weeds in a field, the field comprising weeds and plants of the invention or the herbicide resistant  
30 progeny thereof, the method comprising application to the field of a glyphosate type herbicide in an amount sufficient to control the weeds without substantially affecting the plants. According to this method, one or more of a herbicide, insecticide, fungicide,

nematicide, bacteriocide and an anti-viral may be applied to the field (and thus the plants contained within it) either before or after application of the glyphosate herbicide.

The invention still further provides a method of producing plants which are substantially tolerant or substantially resistant to glyphosate herbicide, comprising the steps  
5 of:

- (i) transforming plant material with the polynucleotide or vector of the invention;
- (ii) selecting the thus transformed material; and
- (iii) regenerating the thus selected material into morphologically normal fertile whole plants.

10 The transformation may involve the introduction of the polynucleotide into the material by any known means, but in particular by: (i) biolistic bombardment of the material with particles coated with the polynucleotide; (ii) by impalement of the material on silicon carbide fibres which are coated with a solution comprising the polynucleotide; or (iii) by introduction of the polynucleotide or vector into *Agrobacterium* and co-cultivation of the  
15 thus transformed *Agrobacterium* with plant material which is thereby transformed and is subsequently regenerated. Plant transformation, selection and regeneration techniques, which may require routine modification in respect of a particular plant species, are well known to the skilled man. The thus transformed plant material may be selected by its resistance to glyphosate.

20 The invention still further provides the use of the present inventive polynucleotide or vector in the production of plant tissues and/or morphologically normal fertile whole plants which are substantially tolerant or substantially resistant to glyphosate herbicide.

The invention still further includes a method of selecting biological material transformed so as to express a gene of interest, wherein the transformed material comprises  
25 the polynucleotide or vector of the invention, and wherein the selection comprises exposing the transformed material to glyphosate or a salt thereof, and selecting surviving material. The said material may be of plant origin, and may in particular be derived from a monocot selected from the group consisting of barley, wheat, corn, rice, oats, rye, sorghum, pineapple, sugar cane, banana, onion, asparagus and leek.

30 The invention still further includes a method for regenerating a fertile transformed plant to contain foreign DNA comprising the steps of:

- (a) producing regenerable tissue from said plant to be transformed;

- (b) transforming said regenerable tissue with said foreign DNA, wherein said foreign DNA comprises a selectable DNA sequence, wherein said sequence functions in a regenerable tissue as a selection device;
- (c) between about one day to about 60 days after step (b), placing said  
5 regenerable tissue from step (b) in a medium capable of producing shoots from said tissue, wherein said medium further contains a compound used to select regenerable tissue containing said selectable DNA sequence to allow identification or selection of the transformed regenerated tissue;
- (d) after at least one shoot has formed from the selected tissue of step (c)  
10 transferring said shoot to a second medium capable of producing roots from said shoot to produce a plantlet, wherein the second medium optionally contains the said compound; and
- (e) growing said plantlet into a fertile transgenic plant wherein the foreign DNA is transmitted to progeny plants in Mendelian fashion, characterised in that the foreign DNA  
15 is, or the selectable DNA sequence comprised by the foreign DNA comprises, the polynucleotide according to any one of claims 1 to 34, and the said compound is glyphosate or a salt thereof. The plant may be a monocot as indicated above - more preferably selected from banana, wheat, rice, corn and barley and the said regenerable tissue may consist of embryogenic calli, somatic embryos, immature embryos etc.

20 The present invention will be further apparent from the following description taken in conjunction with the associated drawings and sequence listings.

### List of Sequences

- SEQ ID NO. 1-40 PCR primers.
- SEQ ID NO. 41 Rice genomic EPSPS sequence (from ATG).
- 25 SEQ ID NO. 42 Rice genomic EPSPS sequence containing double mutation.
- SEQ ID NO. 43 Maize polyubiquitin enhancer.
- SEQ ID NO. 44 Rice actin enhancer 1.
- SEQ ID NO. 45 Rice Genomic G1 EPSPS (to ATG)
- SEQ ID NO. 46 Rice Genomic G3 EPSPS (to ATG).
- 30 SEQ ID NO. 47 Rice Genomic G2 EPSPS +Maize Adh1 intron
- SEQ ID NO. 48 Maize Adh1 intron

**List of Figures**

Figure 1 Rice EPSPS genomic schematic map.

Figure 2 Vector pCR4-OSEPSPS (rice dmEPSPS gene in vector pCR4-Blunt)

Figure 3 Schematic representation of strategy used to introduce the double mutation.

5 Figure 4 Vector pTCV1001

Figure 5 Vector pTCV1001OSEPSPS (comprising rice dmEPSPS gene in vector pTCV1001).

Figure 6 Vector pTCV1001EPSPSPAC (comprising rice dmEPSPS gene in vector pTCV1001).

10 Figure 7 Vector pBluSK+EPSPS (comprising rice dmEPSPS gene in vector pBluescript SK+).

Figure 8 Vector pPAC1

Figure 9 Vector pTCVEPSPSPH

Figure 10 Vector pTCVEPSPSADH

15 Figure 11 Vector pBluSKEPSPSADH (comprising rice dmEPSPS gene containing Adh1 intron)

Figure 12 Vector pIGPD9

Figure 13 Schematic diagram relating to the use of "minimal EPSPS promoters"

Figure 14 Vector Zen 8

20 Figure 15 Vector Zen 19

Figure 16 Vector Zen 21

Figure 17 Introduction of Zen vectors into superbinary vectors

**Production of plants tolerant to glyphosate treatment through over-expression of a mutated EPSPS under the control of a non-heterologous promoter.**

25 The term 'enhancer' as used throughout this specification refers to sequences upstream of a promoter which do not comprise the promoter itself but which act to enhance and regulate transcription from the promoter. The term "EPSPS promoter deletion" as used throughout this patent specification refers to the EPSPS promoter together with nucleotides constituting at least a part of the EPSPS genes native enhancer, ie, EPSPS derived sequences  
30 upstream (5' ) of the EPSPS promoter.

In respect of the transformation of plant material, those skilled in the art will recognise that although particular types of target material (e.g. embryogenic cell suspension

culture or dedifferentiating immature embryos) and particular methods of transformation (e.g. using *Agrobacterium* or particle bombardment) are specified in the examples below, the present invention is not limited to these particular embodiments and such target materials and methods may be used interchangeably. Furthermore, the term "plant cells" as used  
5 throughout this description of the invention can refer to isolated cells, including suspension cultures as well as to cells in an intact or partly intact tissue such as embryo, scutella, microspore, microspore-derived embryo or somatic cells from plant organs. Similarly, although the specific examples are limited to maize, wheat and rice, the invention is equally applicable to any of a broad range of agricultural crops and amenity plants which can be  
10 transformed using suitable methods of plant cell transformation.

General molecular biological methods are carried out according to Sambrook et al (1989) 'Molecular cloning: A laboratory Manual, 2nd Edn. Cold Spring Harbour Lab. Press.

**EXAMPLE 1. Generation of a cDNA probe for Rice EPSPS**

A partial length cDNA encoding rice EPSPS is obtained using reverse transcriptase  
15 PCR (RT-PCR). Total RNA is isolated from two-week-old rice plants (*Oryza sativa* L. indica var. Koshihikari) using the TRI-ZOL™ method (Life Technologies). First-strand cDNA synthesis is performed using Superscript II reverse transcriptase (Life Technologies) with 200 ng of EPSPS degenerate reverse 10 primer (SEQ ID NO.1) and 2 µg of total RNA according to the supplied protocols. Second strand synthesis and cDNA amplification by PCR is  
20 performed using EPSPS degenerate primers 10 and 4 (SEQ ID NO.1 and SEQ ID NO.2) and PCR beads (Pharmacia) according to the manufacturers instructions. All letter codes are standard abbreviations (Eur. J. Biochem. (1985) 150:15)

SEQ ID NO.1

EPSPS degenerate reverse 10                      5' GCACARGCIGCAAGIGARAAIGCCATIGCCAT 3'

25 SEQ ID NO.2

EPSPS degenerate forward 4                      5' GCWGGAAACWGCMTGCGICCRYTIACIGC 3'

The products are cloned into vector pCR2.1 (Invitrogen) using a TA Cloning kit™ as recommended by the supplier. Plasmid is recovered from selected colonies and the sequence analysed by a process involving computer based homology searches (BLAST) to confirm that  
30 the cloned RT-PCR product shows high homology to known plant EPSPS sequences.

## **EXAMPLE 2. Isolation of Rice EPSPS genomic sequence and cloning of the rice EPSPS gene**

A region of genomic DNA containing the full rice EPSPS gene and 5' upstream region is isolated from a  $\lambda$  EMBLSP6/T7 genomic library constructed from five-day-old etiolated rice shoots (*Oryza sativa* L. Indica var. IR36) (Clontech).  $1 \times 10^6$  plaque forming units (pfu) are screened using the  $^{32}\text{P}$ -labelled rice EPSPS cDNA probe (example 1) using protocols provided by the manufacturer. Positive plaques are subjected to subsequent rounds of hybridisation screening until plaque purity of a cross-hybridising plaque is obtained.  $\lambda$ -DNA is prepared from the phage pure stock, according to the method described by Sambrook et al., 1989. The DNA obtained is analysed by restriction digest and Southern blotting, using the same  $^{32}\text{P}$ -labelled rice EPSPS cDNA as a probe. Restriction fragments that cross-hybridise are, where applicable, blunt-ended using a method such as Perfectly Blunt™ (Novagen), and cloned into a suitable vector such as pSTBlue (Novagen). The DNA is then sequenced using an ABI 377A PRISM automated DNA sequencer. Figure 1 shows a schematic of the rice EPSPS gene with some of the restriction sites marked.

A 3.86 kb fragment of the rice EPSPS gene, containing the coding region, the EPSPS promoter, some of the 5' upstream region and the terminator is obtained by PCR. Oligonucleotide primer OSGRA1 (SEQ ID NO.3) is used in conjunction with OSEPSPS3 (SEQ ID NO. 4) to amplify the desired region. OSEPSPS3 contains additional *Sac* I and *Sma* I restriction enzyme sites to facilitate the subcloning of the gene during the later stages of vector construction. A schematic location of these primers is given in Figure 1.

### **SEQ ID NO. 3**

OSSGRA1 5' ATTCTTCTTCTTCTCCTCCCTTCTCCGCCTC 3'

### **SEQ ID NO. 4**

OSEPSPS3 5' GAGCTCCCCGGGCGAGTGTGTTGTGTTCTGTCTAATG 3'

High fidelity Pfu Turbo™ polymerase (Stratagene) is used to perform the PCR reaction with DNA obtained from  $\lambda$  preparation (described above) as the amplification template. The PCR product of expected size is cloned into pCRblunt 4-TOPO™ (Invitrogen) and sequenced to check integrity.

## **EXAMPLE 3. Mutation of T to I and P to S in the rice EPSPS.**

The T to I and P to S mutation is obtained by the introduction of two point mutations. These mutations are introduced into the rice genomic EPSPS gene by PCR using

oligonucleotide primers containing the desired mutation. A schematic diagram, indicating the binding sites of the primers used, is shown in Figure 3. Two separate PCR reactions are performed (both using the  $\lambda$  DNA as template).

1) EcoRVEnd (SEQ ID NO.5) + OSMutBot (SEQ ID NO. 6)

5 2) OsMutTop (SEQ ID NO. 7) + SalIEnd (SEQ ID NO. 8)

SEQ ID NO.5

EcoRVEnd 5' GCTTACGAAGGTATGATATCCTCCTACATGTCAGGC 3'

SEQ ID NO.6

OSMutBot 5' GCAGTCACGGCTGCTGTCAATGATCGCATTGCAATTCCAGCGTTCC 3'

10 SEQ ID NO.7

OsMutTop 5' GGAACGCTGGAATTGCAATGCGATCATTGACAGCAGCCGTGACTGC 3'

SEQ ID NO.8

SalIEnd 5' GGTGGGCATTTCAGTGCCAAGGAAACAGTCGACATCCGCACCAAGTTGTTTCAACC 3'

The resulting PCR products are joined by using equimolar concentrations of each  
15 PCR product as template with the two oligos SalIEnd and EcoRVEnd in a new PCR reaction. An aliquot of the reaction product is analysed by agarose gel electrophoresis and cloned into pCR-Blunt II™ (Invitrogen). Plasmid DNA is recovered and sequenced to detect the successful incorporation of the double mutation.

The DNA fragment containing the double mutation is incorporated into the rice  
20 EPSPS genomic clone (Figure 1) as follows. The clone containing the double mutant is digested with *Eco* RV and *Sal* I. The plasmid containing the rice EPSPS DNA PCR product is similarly digested and the *Eco* RV/*Sal* I fragment containing the double mutant ligated into the rice EPSPS gene in pCR4Blunt -TOPO™ using standard cloning methods described in Sambrook *et al.*, 1989 and transformed into competent *E. coli*. Plasmid is recovered from  
25 resultant colonies and sequenced in order to confirm the presence of the double mutation with no further alterations. This plasmid, pCR4-OSEPSPS, is shown in Figure 2. The genomic rice EPSPS gene containing the double mutant (Figure 2) is excised from pCR4-Blunt-TOPO™ using *Pst* I and *Not* I and ligated into vector pTCV1001 (Figure 4), to generate pTCV10010SEPSPS (Figure 5) and this is transformed into *E. coli* for  
30 amplification. Next, the *Pac* I/*Eco* RV restriction fragment is excised from the  $\lambda$  DNA (figure 1) and inserted into pTCV10010SEPSPS (figure 5) to generate pTCV1001EPSPSPAC (Figure 6). The rice dmEPSPS gene, now containing sequence from

*Pac* I to *Sac* I (Figure 6), is excised from pTCV1001EPSPSPAC (Figure 6) as an *Eag* I/*Sac* I fragment and ligated into similarly digested pBluescript SK+ to make pBluSK+EPSPS (Figure 7). Further rice EPSPS upstream regions and desired enhancers are assembled (as described below) and ligated into the pBluescript SK+ vector using *Xba* I/*Pac* I.

5 **EXAMPLE 4. Generation of singly enhanced : rice EPSPS promoter fusions**

Figure 1 indicates the binding sites of the primers G1 and G2 used to generate a series of deletions at the 5' end of the rice EPSPS gene. The G1 and G2 primers (SEQ ID NO 9 and SEQ ID NO 10) are used in combination with the RQCR10 primer (SEQ ID NO 11) using the rice EPSPS lambda DNA template and Pfu Turbo™ polymerase (Stratagene) using  
10 protocols provided by the supplier.

SEQ ID NO.9

G1                    5' CGCCTGCAGCTCGAGGTTGGTTGGTGAGAGTGAGACACC 3'

SEQ ID NO.10

G2                    5' CGCCTGCAGCTCGAGGCCACACCAATCCAGCTGGTGTGG 3'

15 SEQ ID NO.11

RQCR10            5' GAACCTCAGTTATATCTCATCG 3'

The products obtained are analysed by agarose gel electrophoresis and cloned into pCR-Blunt II-TOPO™ vector (Invitrogen). The sequence of the resulting products is determined to ensure that there is no alteration in the sequence of the rice genomic EPSPS  
20 clone. Clones to progress are selected based on their orientation within the vector by establishing whether or not *Xho* I digestion removes only the polylinker sequence rather than the whole insert from the vector.

The sequence of the maize polyubiquitin and rice actin genes and their associated 5' upstream regions are published in the EMBL database (U29159 and X15865 respectively).  
25 Primers are designed so as to amplify only the upstream enhancer regions of the said genes. The maize polyubiquitin enhancer (SEQ ID NO. 43) is thus obtained by PCR using primers SEQ ID NO. 12 and SEQ ID NO. 13 in conjunction with Pfu Turbo™ polymerase and maize genomic DNA as the template. These primers both contain a *Spe* I restriction site to facilitate further manipulations of the enhancer (note, however, that the *Xho* I site present within the  
30 maize polyubiquitin enhancer is utilised as the 3' restriction site). The rice actin enhancer (SEQ ID NO. 44) is obtained in a similar manner using primers (SEQ ID No 14 and SEQ ID



No 15) with rice genomic DNA as template. These primers contain a *Xba* 1 and *Pst* 1 restriction site respectively to facilitate further manipulations of the enhancer.

The following oligonucleotide primers are used.

SEQ ID NO.12

5 MPU5 5' GCGGCCGCACTAGTGGCCGCCATCAGCGCCAGCTTTTGTTC 3'

SEQ ID NO.13

MPU3 5' TTAAGTAGTGAGGAGGCCGCTGCGTGC 3'

SEQ ID NO.14

RA5 5' CGCCTCTAGAGGCCGCCGATATCCCTCAGCCGCCTTTCACTATC 3'

10 SEQ ID NO.15

RA3 5' CGCTGCAGTGCTCGCGATCCTCCTCGCTTTTCC 3'

The sequence of the amplified and cloned molecules is confirmed following cloning into the PCR Blunt-II-TOPO vector (Invitrogen). The pCR Blunt\_II-TOPO vector, containing the EPSPS 5'UTR deletion is digested with either *Not* 1/*Xho* 1 (MPU) or *Xba* 1/*Pst* 1 (RA). The Enhancer is removed from its respective pCR Blunt-II-TOPO vector also using required restriction enzymes and ligated into the first vector containing the 5'UTR EPSPS deletion.

#### **EXAMPLE 5. Generation of doubly enhanced : rice EPSPS promoter fusions.**

In order to further increase expression from the rice EPSPS promoter a second rice actin enhancer is incorporated into the existing rice actin:EPSPS fusion. To achieve this end, enhancer/EPSPS fusions are made initially(as described in example 4) comprising a single (first) rice actin enhancer. The second rice actin enhancer is amplified using the primers RAPST (SEQ. ID. NO. 16) and RAPAC (SEQ ID NO 17). These primers facilitate the introduction of a *PST* 1 site at the 5' terminus and a *Pac* 1 site at the 3' terminus of the enhancer.

SEQ ID NO.16

RAPST 5' gcgctgcagGATATCCCTCAGCCGCCTTTCACTATC 3'

SEQ ID NO.17

RAPAC 5' gcgttaattaaTGCTCGCGATCCTCCTCGCTTTTCC 3'

30 Once sequenced, the PCR product (as *Pst* 1 : *Pac* 1) is introduced into the construct which comprises the first rice actin enhancer : G1 EPSPS gene fusion (example 4).

**EXAMPLE 6. Insertion of Adh1 intron into the 5' UTR of the rice EPSPS gene**

The insertion of the Maize Adh1 intron 1 into the desired rice EPSPS promoter deletion (e.g. made as described in Example 4) is performed prior to the generation of the fusion construct with the desired enhancer(s). In this particular example the Adh1 intron is introduced into the G2 EPSPS promoter deletion. The skilled man will appreciate that similar methodology can be adopted to incorporate the Adh1 intron into other EPSPS promoter deletions. The maize Adh1 intron is inserted into the constructs by PCR. The Adh 1 intron is amplified from a suitable source, such as maize genomic DNA or a vector such as pPAC1 (Figure 8) using primers Adh5 (SEQ ID NO. 18) and Adh3 (SEQ ID NO. 19):

SEQ ID NO. 18

Adh5 cccatcctccccgacctccacgccgcggcaggatcaagtgcaaagggtccgccttgtttctctctctg

SEQ ID NO. 19

Adh3 gacgccatgggtcgccgccatccgcagctgcacgggtccaggaaagcaatc

The resulting PCR product is denatured and used as a primer in conjunction with Adh5Pac (SEQ ID NO. 20) to amplify the desired product using the vector pTCV1001EPSPSPAC (Figure 2) as template.

SEQ ID NO. 20

Adh5Pac cgagttcttatagtagatttcaccttaattaaaac

The resulting PCR product is cloned into PCR-blunt II (Invitrogen). The *Pac* 1:*Hind* III fragment is excised from the rice genomic clone (Figure 1) and inserted into pTCV1001 to generate pTCVEPSPSPH (Figure 9). Next, the *Pac* I/*Nco* I PCR product comprising the Adh1 intron is inserted into pTCVEPSPSPH as shown in the schematic (Figure 9). The *Pac* 1:*Eco* RV fragment present in the cloned EPSPS gene containing the double mutant (Figure 10) is excised and replaced with the *Pac* I/*Eco* RV fragment from pTCVEPSPSPH that comprises the Adh1 intron sequences (Figure 9). Finally the full EPSPS gene comprising the Adh 1 sequence is excised from pTCVEPSPSPH as an *Eag* I/*Sac* I fragment and cloned into pBluescript SK+ to give pBluSKEPSPSADH (figure 11).

**EXAMPLE 7. Introduction of optimised pre ATG consensus sequence (Kozak) via site directed mutagenesis for constructs comprising the maize adh1 intron.**

Optionally, site directed mutagenesis is performed on constructs containing the Adh1 intron using the QuickChange Site Directed Mutagenesis kit (Stratagene). This is performed on the *Pac*1/*Sac*1 EPSPS fragment in pBluescript SK+ (Figure 11) prior to fusion with the

enhancer : EPSPS promoter fusions. The following oligonucleotides are used according to the supplied protocols to optimise the KOZAK sequence.

SEQ ID NO. 21

Oskozak 5' GGACCCGTGCAGCTGCGGTACCATGGCGGCGACCATGGC 3'

5 SEQ ID NO. 22

OSkozakrev 5' GCCATGGTCGCCGCCATGGTACCGCAGCTGCACGGGTCC 3'

Clones are analysed by restriction analysis, using *Kpn* 1, on recovered plasmid. The correctly altered DNA is characterised by an additional *Kpn* 1 restriction site compared to the un-altered DNA. The sequence is then verified by automated DNA sequencing.

10 The altered DNA sequence may be transferred original constructs using the unique restriction enzyme sites of *Sph* 1 or *Pac* 1 at the 5' end and *Avr* II or *Eco* RV at the 3' end as appropriate for each vector.

**EXAMPLE 8. Completion of EPSPS expression cassettes comprising, in the 5' to 3'**

15 **direction, Enhancer region(s), rice EPSPS promoter upstream region, EPSPS promoter, EPSPS 5'UTR + (optional) maize Adh1 intron 1, rice EPSPS transit peptide coding region, rice mature EPSPS coding region and rice EPSPS gene terminator region .**

The singly and doubly enhanced rice EPSPS promoter fusions (Examples 4 and 5) contained within the pCR Blunt-II-TOPO vectors are excised using *Xba* 1 and *Pac* 1 (RA) or 20 Not 1 and *Pac* 1 (MPU) and inserted into the similarly digested pBluescript SK+ clone containing the remainder of the rice EPSPS sequence (Figures 7/11). This final cloning step results in the required gene constructs. Examples of constructs (EPSPS expression cassettes) obtainable using the above strategies are given below in Table 1. Schematic maps are given in Figures 14-16.

25

Clone	First enhancer	Second enhancer	EPSPS Promoter deletion	5' UTL Intron	EPSPS genomic coding region	EPSPS Terminator
ZEN6	RA	None	G1	No	Yes	Yes
ZEN10	MPU	None	G1	No	Yes	Yes
ZEN13	RA	RA	G3	No	Yes	Yes
ZEN26	RA	None	Minimal	RA	Yes	Yes

## Optional Further Assembly of DNA constructs

### Use of minimal EPSPS promoters

The promoter region of both the rice actin promoter and the maize polyubiquitin promoters is well defined. In these examples the native promoter of these genes, comprising the "TATA" box, is replaced with that of the rice EPSPS promoter. In this example the EPSPS promoter is used to replace the promoter region in the rice actin gene. The skilled man will appreciate that a similar methodology may be used with a variety of genes. The EPSPS promoter is introduced into the rice actin gene by PCR. Initially, four independent PCR reactions are performed. Primers RA5E (SEQ ID NO. 23) and RA3E (SEQ ID NO. 24) are used with rice genomic DNA template to amplify the rice actin enhancer element; primers RA5I (SEQ ID NO. 25) and RA3I (SEQ ID NO. 26) are used with rice genomic DNA to amplify the rice actin intron; primers EPROM53 (SEQ ID NO. 27) and EPROM3 (SEQ ID NO. 28) are used to amplify the region of rice EPSPS comprising the promoter; and primers REPS5 (SEQ ID NO. 29) and REPS3 (SEQ ID NO. 30) are used to amplify the rice EPSPS gene between the translation start site and the *EcoRV* site (see Figure 1). Each individual PCR product is joined, in order, by successive PCR since each primer used to amplify the region contains a linker to the next. A schematic representation of the process is given in Figure 13.

RA5E            5'tctctagactcagccgcctttcactac3'

(SEQ ID NO. 24)

RA3E            5'aaacccgggttggaagcggaggagGAAGGAGGAGATAAAG 3'

(SEQ ID NO. 25)

RA5I            5' ACCCTCCCCTCTCtaaatcgattggtgggaggaggagag 3'

(SEQ ID NO. 26)

RA3I            5' ggtctacctacaaaaaagctccgcacgagGGTACCGCCGCTGGTAC 3'

(SEQ ID NO. 27)

EPROM53      5' CCTTCGCCTCCCCTCcttctctctatttcttc 3'

(SEQ ID NO. 28)

EPROM3        5' gttggtgggaggaggagATTTAGCTAACCACC

(SEQ ID NO. 29)

REPS5          5' GTTTTTTCGAGGCGTGCTCccatggcggcgacctggcgctcc 3'

(SEQ ID NO. 30)

REPSPS3     5' ggaggatatcataccttcgtaagc 3'

The final DNA fragment obtained, comprising the rice actin enhancer, EPSPS promoter, rice actin intron, and rice EPSPS gene to *Eco* RV site is introduced into pBluSK+EPSPS (Figure 7) as *Xba* 1 / *Eco* RV to give, for example, ZEN26. The complete expression cassette may  
5 then be excised as *Xma* 1 for further subcloning.

The man skilled in the art will appreciate that different lengths of EPSPS promoter can be utilised and that different components, such as the maize polyubiquitin enhancer and intron may be utilised in a similar manner.

**EXAMPLE 10. Preparation of DNA for plant transformation**

10         The above procedure describes the assembly of 'EPSPS expression cassettes' comprising, in a 5' to 3' direction, an enhancer sequence(s), an EPSPS promoter from rice, a region encoding a rice EPSPS transit peptide, a region encoding a mature rice EPSPS enzyme which is resistant to glyphosate through having T to I and P to S changes at the specified positions and a rice EPSPS gene terminator.

15         Optionally the desired cassettes also further comprise a drug selection marker gene (e.g. ampicillin resistance, kanamycin resistance etc.) a T-DNA Left or Right Border region and (optionally) a scaffold attachment region added 5' and/or 3' to the above described construct. The skilled man will recognise that similar methods to those described above can be used to obtain these added components and clone them into the desired positions.

20         **EXAMPLE 11. Transformation of corn lines using an *Agrobacterium* strain containing a superbinary vector which includes an EPSPS expression cassette between the right and left borders of the T-DNA; selection and regeneration of plant cells and plants which are resistant to glyphosate**

**Construction of *Agrobacterium* strain**

25         Bluescript plasmid DNA (e.g. ZEN 7, 8, 17, 19, 21 and 22) is digested with either *Xma* 1 or with *Xba* 1 / *Sac* 1 and the thus-obtained (~ 5.5- 7 kb) EPSPS-encoding fragment ligated into a position within the cloning site located between the right and left T-DNA borders of similarly restricted pSB1. In the case, for example, of using the *Xma* 1 fragment of pZEN 8 this ligation creates the plasmid pZEN8SB11 (Figure 16). The construction of  
30 plasmid pSB11 and the construction of its parent, pSB21, is described by Komari et al (1996, Plant J. 10: 165-174). The T-DNA region of pZEN8 is integrated into the superbinary pSB1 vector.(Saito *et al* EP 672 752 A1) by a process of homologous recombination (Figure 17) to

create the plasmid, pSB1ZEN8. To achieve this the plasmid pZEN8SB11 is transformed into *E. coli* strain HB101 which is then, according to the triple cross method of Ditta et al (1980, Proc. Natl. Acad. Sci. USA 77: 7347-7351), mated with an *Agrobacterium* LBA4404 harbouring pSB1 to create the transformed strain of *Agrobacterium*, LBA4404 (pSB1ZEN8) in which the presence of the cointegrate plasmid pSB1ZEN8 is selected for on the basis of resistance to spectinomycin. The identity of pSB1ZEN8 is also confirmed on the basis of Sal I restriction analysis (Figure 17). LBA4404 strains containing the directly analogous constructs pSB1ZEN7, pSB1ZEN17, pSB1ZEN19, pSBZEN21 and pSB1ZEN22 are similarly constructed starting from the XmaI fragments of pZEN7, ZEN17, ZEN19, ZEN21 and ZEN22.

Alternatively, using similar methods to those described above, a similar fragment of pZEN7, ZEN 8 etc is homologously recombined into a position between the right and left borders of the superbinary vector pTOK162 (Fig 1 in US 5591616) to generate a similar set of cointegrate plasmids selected for in *Agrobacterium* on the basis of combined resistance to kanamycin and spectinomycin.

*Agrobacterium* strain LBA4404 which has a helper plasmid PAL4404 (having a complete *vir* region) is available from the American Type Culture Collection (ATCC 37349). An alternative useful strain is *Agrobacterium* EHA101 (1986, Hood et al, J. Bacteriol., 168(3): 1283-1290) which has a helper plasmid having the *vir* region from the strongly virulent strain *Agrobacterium tumefaciens* A281.

#### Preparation of *Agrobacterium* suspensions

*Agrobacterium* strains LBA4404(pSB1ZEN7), LBA4404 (pSB1ZEN8) etc are each streaked onto plates containing 'PHI-L' solid medium and cultured at 28 C in the dark for 3 to 10 days.

PHI-L medium is as described on page 26 (Example 4) of WO 98/32326. PHI-L medium made up in double-distilled water comprises 25 ml/l of stock solution A, 25 ml/l of stock solution B, 450.9 ml/l of stock solution C and 50 mg/l of spectinomycin. Stock solutions are sterilised by autoclaving or filtration. Stock solution A is 60 g/l  $K_2HPO_4$  and 20 g/l  $NaH_2PO_4$  adjusted to pH 7.0 with KOH: stock solution B is 6 g/l  $Mg SO_4 \cdot 7H_2O$ , 3 g/l KCl, 20 g/l  $NH_4Cl$ , 0.2 g/l  $CaCl_2$  and 50 mg/l  $FeSO_4 \cdot 7H_2O$ : stock solution C is 5.56 g/l of glucose and 16.67 g/l of agar (A-7049, Sigma Chemicals, St Louis, Mo, USA)

Alternatively the *Agrobacterium* are cultured for 3 -10 d on a plate containing YP medium (5 g/l yeast extract, 10 g/l peptone, 5 g/l NaCl, 15 g/l agar at pH 6.8) as described by Ishida et al (1996, Nature Biotechnology, 14, 745-750) or, alternatively, as described by Hei et al in US 5591616 (AB medium (Drlica and Kado, 1974; Proc. Natl. Acad. Sci. USA 71:3677-3681)) but, in each case, modified to provide the appropriate antibiotic selection (e.g. containing 50 mg/ ml spectinomycin in the case of *Agrobacterium* strain LBA4404(pSB1ZEN7) etc. or containing both 50 mg/ ml spectinomycin and 50 mg/ ml kanamycin in the case that *Agrobacterium* containing a pTOK 162-derived superbinary vector is used).

Plates of *Agrobacterium* made as described above are stored at 4 C and used within a month of preparation. For preparation of suspensions a single colony from the master plate is streaked out onto a plate containing, at pH 6.8, 5 g/l yeast extract (Difco), 10 g/l peptone (Difco), 5 g/l NaCl, 15 g/l agar (Difco) and 50 mg/l of spectinomycin (or as appropriate for the particular strain of *Agrobacterium*) . Plates are incubated at 28 C, in the dark for 2d.

Suspensions of *Agrobacterium* for transformation of plant material are prepared in a similar manner to described in US 5591616. (Using good microbiological practice to avoid contamination of aseptic cultures) 3 X 5 mm loopfuls of *Agrobacterium* are removed from plates, transferred and suspended in 5 ml of sterile AA liquid medium in a 14 ml Falcon tube. As used here, AA liquid medium at pH 5.2 contains the major inorganic salts, amino acids and vitamins defined by Toriyama and Hinata (1985) in Plant Science 41, 179-183), the minor inorganic salts of Murashige and Skoog medium (Murashige and Skoog, 1962 in Physiol. Plant 15, 473-497), 0.5 g/l of casamino acids (casein hydrolysate), 1 mg/l of 2,4 - dichlorophenoxyacetic acid (2,4-D), 0.2 mg/l of kinetin, 0.1 mg/l of gibberellin, 0.2M glucose, 0.2M sucrose and 0.1 mM acetosyringone.

Alternatively, suspensions of *Agrobacterium* for transformation of plant material are prepared in a similar manner to described in WO 98/32326. 3 X 5 mm loopfuls of *Agrobacterium* are removed from plates, transferred and suspended in 5 ml of the sterile PHI-A basic medium as described in Example 4 on page 26 of WO 98/32326 or, alternatively, suspended in 5 ml of the sterile PHI-I combined medium also described in Example 4 on page 26 of WO 98/32326. In either case 5 ml of 100 mM 3'-5'-Dimethoxy-4'-hydroxyacetophenone is also added. PHI-A basic medium at pH 5.2 comprises 4 g/l of CHU(N6) basal salts (Sigma C-1416), 1.0 ml/l of Eriksson's vitamin mix (1000X , Sigma E-

1511), 0.5 mg/l thiamine. HCl, 1.5 mg/ ml of 2,4-D, 0.69g/ l L-proline, 68.5 g/ l sucrose and 68.5 g/ l glucose. PHI-I combined medium, also adjusted to pH 5.2 with KOH and filter sterilized, comprises 4.3 g/l of MS salts (GIBCO-BRL), 0.5 mg/ ml nicotinic acid, 0.5 mg/ ml pyridoxine. HCl, 1.0 mg/ ml thiamine. HCL, 100 mg/ l myo-inositol, 1 g/ l vitamin assay casamino acids (Difco), 1.5 mg/ ml of 2,4-D, 0.69g/ l L-proline, 68.5 g/ l sucrose and 36 g/ l glucose.

Alternatively, suspensions of *Agrobacterium* for transformation of plant material are prepared in a similar manner to described by Ishida et al (1996) Nature Biotechnology, 14, 745-750. 3 X 5 mm loopfuls of *Agrobacterium* are removed from plates, transferred and suspended in 5 ml of LS-inf medium. LS-inf medium (Linsmaier and Skoog, 1965, Physiol. Plant 18, 100-127) adjusted to pH 5.2 with KOH contained LS major and minor inorganic salts, 0.5 mg/ ml nicotinic acid, 0.5 mg/ ml pyridoxine. HCl, 1.0 mg/ ml thiamine. HCL, 100 mg/ l myo-inositol, 1 g/ l vitamin assay casamino acids (Difco), 1.5 mg/ ml of 2,4-D, 68.5 g/ l sucrose and 36 g/ l glucose.

However produced, the suspension of *Agrobacterium* is vortexed to make an even suspension and the cell population adjusted to between  $0.5 \times 10^9$  and  $2 \times 10^9$  cfu/ ml (preferably the lower).  $1 \times 10^9$  cfu/ ml corresponds to an OD (1 cm) of ~ 0.72 at 550 nm.

*Agrobacterium* suspensions are aliquoted into 1 ml lots in sterile 2 ml microcentrifuge tubes and used as soon as possible

#### 20 Corn lines for transformation

Suitable maize lines for transformation include but are not restricted to, A188, F1 P3732, F1 (A188 x B73Ht), F1 (B73Ht x A188), F1 (A188 x BMS). Varieties A188, BMS (Black Mexican Sweet) and B73 Ht are obtained from the Ministry of Agriculture, Forestry and Fisheries. P3732 is obtained from IWATA RAKUNOU KYODOKUMIAI. Suitable maize lines also include a variety of A188 x inbred crosses (e.g PHJ90 x A188, PHN46 x A188, PHPP8 x A188 in table 8 of WO98/ 32326) as well as elite inbreds from different heterotic groups (e.g PHN46, PHP28 and PHJ90 in table 9 of WO98/ 32326).

For example immature embryos are produced from "Hi-II" corn. "Hi-II" is a hybrid between inbreds (A188 x B73) generated by reciprocal crosses between Hi-II parent A and Hi-II parent B available from the Maize Genetic Cooperation Stock Center, University of Illinois at Champaign, Urbana, Illinois). Seeds, termed 'Hi-II' seeds obtained from these



crosses are planted out in a greenhouse or field. The resulting Hi-II plants are self or cross-pollinated with sister plants

Preparation of immature embryos, infection and co-cultivation

Transformation of immature embryos of corn is carried out by contacting the  
5 immature embryos with the suitable recombinant strains of *Agrobacterium* described above. An immature embryo means the embryo of an immature seed which is in the stage of maturing following pollination. Immature embryos are an intact tissue that is capable of cell division to give rise to callus cells that can then differentiate to produce the tissues and organs of a whole plant. Preferred material for transformation also includes the scutella of  
10 embryos which is also capable of inducing dedifferentiated calli with the ability to regenerate normal fertile plants having been initially transformed. Preferred material for transformation thus also includes callus derived from such dedifferentiated immature zygotic embryos or scutella.

Immature corn embryos are isolated aseptically from developing ears as described by  
15 Green and Phillips (1976, Crop. Sci. 15: 417-421) or, alternatively, by the methods of Neuffer et al (1982, "Growing Maize for genetic purposes" in *Maize for biological research*, W.F. Sheridan ed., University Press, University of North Dakota, Grand Forks, North Dakota, USA). For example, immature corn embryos between 1-2 mm (preferably 1-1.2 mm) long are aseptically isolated from female spikes at 9-12 (preferably 11) d after  
20 pollination using a sterile spatula. Typically ears are surface sterilised with 2.63% sodium hypochlorite for 20 min before washing with sterile deionized water and aseptic removal of immature embryos. Immature embryos (preferably ~ 100 in number) are dropped directly into a 2 ml microcentrifuge tube containing about 2 ml of the same medium as used for preparing the suspension of *Agrobacterium* (the alternatives for which are described above).  
25 The cap of the tube is closed and the contents mixed by vortexing for a few seconds. The medium is decanted off, 2 ml of fresh medium are added and vortexing is repeated. All of the medium is then drawn off to leave the washed immature embryos at the bottom of the tube.

Having prepared the immature maize embryos the next phase of the process, the  
30 infection step, is to contact them in with the transformed strain of *Agrobacterium*.

In one example of this process, the infection step takes place in a liquid medium which includes the major inorganic salts and vitamins of N6 medium (1987, Chu C.C. Proc.

Symp. Plant Tissue Culture, Science Press Peking. Pp 43-50) as described in example 4 of WO 98/32326. 1.0 ml of suspension of *Agrobacterium*, prepared as described above in PHI-A medium is added to the embryos in the microcentrifuge tube and vortexed for about 30s. Alternatively, 1.0 ml of suspension of *Agrobacterium* prepared, also as described above, in either PHI-I medium or in LS-inf medium is added.

After standing for 5 minutes the suspension of *Agrobacterium* and embryos is poured out into a Petri plate containing either 1) PHI-B medium or 2) PHI-J medium or 3) LS-AS medium according to whether the original suspension of *Agrobacterium* had been prepared in PHI-A medium, PHI-I medium or LS-inf medium, respectively. The *Agrobacterium* suspension is drawn off using a Pasteur pipette, the embryos manipulated so that they sit axis-side downwards onto the medium, the plate sealed with parafilm and incubated in the dark at 23-25 C for 3 days of cocultivation. PHI-B medium at pH 5.8 comprises 4 g/l of CHU(N6) basal salts (Sigma C-1416), 1.0 ml/l of Eriksson's vitamin mix (1000X, Sigma E-1511), 0.5 mg/l thiamine. HCl, 1.5 mg/ml of 2,4-D, 0.69g/l L-proline, 0.85 mg/l silver nitrate, 30 g/l sucrose, 100 mM acetosyringone and 3 g/l gelrite (Sigma). PHI-J medium, also adjusted to pH 5.8 comprises 4.3 g/l of MS salts (GIBCO-BRL), 0.5 mg/ml nicotinic acid, 0.5 mg/ml pyridoxine. HCl, 1.0 mg/ml thiamine. HCL, 100 mg/l myo-inositol, 1.5 mg/ml of 2,4-D, 0.69g/l L-proline, 20 g/l sucrose, 10 g/l glucose, 0.5 g/l MES (Sigma), 100 mM acetosyringone and 8 g/l purified agar (Sigma A-7049). LS-AS medium (Linsmaier and Skoog, 1965, Physiol. Plant 18, 100-127) adjusted to pH 5.8 with KOH contains LS major and minor inorganic salts, 0.5 mg/ml nicotinic acid, 0.5 mg/ml pyridoxine. HCl, 1.0 mg/ml thiamine. HCL, 700 mg/l L-proline, 100 mg/l myo-inositol, 1.5 mg/ml of 2,4-D, 20 g/l sucrose, 10 g/l glucose, 0.5 g/l MES, 100 mM acetosyringone and 8 g/l purified agar (Sigma A-7049).

Following the preparation of immature embryos, as described above, an alternative method of achieving transformation is to infect them during and after a period of dedifferentiation as described in US 5591616. Immature embryos are placed on LSD 1.5 solid medium containing LS inorganic salts and vitamins along with 100 mg/ml casamino acids, 700 mg/l L-proline, 100 mg/l myo-inositol, 1.5 mg/ml of 2,4-D, 20 g/l sucrose and 2.3 g/l of gelrite. After 3 weeks at 25 C, calli originating from the scutella are collected in a 2 ml microcentrifuge tube and immersed in 1 ml of *Agrobacterium* suspension prepared, as described above, in AA medium. After standing for 5 minutes, the resultant calli are

transferred to 2N6 solid medium containing 100  $\mu$ M acetosyringone and incubated in the dark at 25 C for a 3 day period of cocultivation. 2N6 solid medium comprises the inorganic salts and vitamins of N6 medium (Chu C.C., 1978; Proc. Symp. Plant Tissue Culture, Science Press Peking, pp 43-50) containing 1 g/l casamino acids, 2 mg/l 2,4-D, 30 g/l sucrose and 2 g/l of gelrite.

'Resting and Selection of transformants'

Following cocultivation, embryos are, optionally, transferred to a plate containing PHI-C medium, sealed over with parafilm and incubated in the dark for 3 days for a 'resting step' prior to selection. PHI-C medium at pH 5.8 comprises 4 g/l of CHU(N6) basal salts (Sigma C-1416), 1.0 ml/l of Eriksson's vitamin mix (1000X, Sigma E-1511), 0.5 mg/l thiamine. HCl, 1.5 mg/ml of 2,4-D, 0.69g/l L-proline, 0.85 mg/l silver nitrate, 30 g/l sucrose, 0.5 g/l MES, 100 mg/l carbenicillin and 8 g/l purified agar (Sigma A-7049). As disclosed in WO 98/32326, the desirability of including this resting step in the overall transformation process varies according to corn line and is a matter of experiment.

For the selection step, about 20 embryos are transferred onto each of a number of fresh plates containing PHI-D selection medium or LSD 1.5 selection medium, sealed with parafilm and incubated in the dark at 28 C. PHI-D selection medium, adjusted to pH 5.8 with KOH, comprises 4 g/l of CHU(N6) basal salts (Sigma C-1416), 1.0 ml/l of Eriksson's vitamin mix (1000X, Sigma E-1511), 0.5 mg/l thiamine. HCl, 1.5 mg/ml of 2,4-D, 0.69g/l L-proline, 0.85 mg/l silver nitrate, 30 g/l sucrose, 0.5 g/l MES, 100 mg/l carbenicillin, 8 g/l purified agar (Sigma A-7049) and between 0.1 mM and 20 mM of tissue culture grade N-(Phosphonomethyl)-glycine (Sigma P-9556). LSD 1.5 selection medium, adjusted to pH 5.8 with KOH, comprises LS major and minor inorganic salts (Linsmaier and Skoog, 1965, Physiol. Plant 18, 100-127), 0.5 mg/ml nicotinic acid, 0.5 mg/ml pyridoxine. HCl, 1.0 mg/ml thiamine. HCL, 700 mg/l L-proline, 100 mg/l myo-inositol, 1.5 mg/ml of 2,4-D, 20 g/l sucrose, 0.5 g/l MES, 250 mg/l cefotaxime, 8 g/l purified agar (Sigma A-7049) and between 0.1 mM and 20 mM of tissue culture grade N-(Phosphonomethyl)-glycine (Sigma P-9556).

Alternatively, in the case that the starting material for selection are calli-derived from immature embryos as disclosed in WO 5591616 then such calli are washed with sterilised water containing 250 mg/l cefotaxime before culturing on LSD 1.5 selection medium.

The embryos or clusters of cells that proliferate from the immature embryos are transferred (if necessary using a sterile scalpel) to plates containing fresh selection medium at 2 weekly intervals over a total period of about 2 months. Herbicide-resistant calli are then bulked by continued growth on the same medium until the diameter of the selected callus  
5 exceeds about 1.5 cm

The concentration of N-(Phosphonomethyl)-glycine in the selection medium is chosen appropriately to select a desirable number of genuine transformants and is preferably within the range 0.3- 5 mM. Preferably the concentration of N-(Phosphonomethyl)-glycine used in the selection medium is about 1 mM for the first two weeks of selection and about 3  
10 mM thereafter.

Regeneration of transformants/ propagation and analysis of transformed plant material

The selected calli are regenerated into normal fertile plants according to, for example, the methods described by Duncan et al (1985, Planta, 165, 322-332) by Kamo et al (1985, Bot. Gaz. 146(3), 327-334) and/or by West et al (1993, The Plant Cell, 5, 1361-1369) and/or  
15 by Shillito et al (1989) Bio/ Technol. 7, 581-587.

For example, selected calli of diameter 1.5- 2 cm are transferred to regeneration/ maturation medium and incubated in the dark for about 1-3 weeks to allow the somatic embryos to mature. A suitable regeneration medium, PHI-E medium (WO 98/ 32326) is adjusted to pH 5.6 with KOH and comprises 4.3 g/l of MS salts (GIBCO-BRL), 0.5 mg/ ml nicotinic acid,  
20 0.5 mg/ ml pyridoxine. HCl, 0.1 mg/ ml thiamine. HCL, 100 mg/ l myo-inositol, 2 mg/ l glycine, 0.5 mg/ l zeatin, 1.0 mg/ ml of indoleacetic acid, 0.1 mM abscisic acid, 100 mg/ l carbenicillin, 60 g/ l sucrose, 8 g/ l purified agar (Sigma A-7049) and, optionally, between 0.02 mM and 1 mM of tissue culture grade N-(Phosphonomethyl)-glycine (Sigma P-9556).

The calli are then transferred to rooting/ regeneration medium and grown at 25 C  
25 under either a schedule of 16 h daylight ( $270 \text{ mE m}^{-2} \text{ s}^{-1}$ ) and 8 h of darkness or under continuous illumination ( $\sim 250 \text{ mE m}^{-2} \text{ s}^{-1}$ ) until such a time as shoots and roots develop. Suitable rooting/ regeneration media are either LSZ medium as described in the following paragraph (optionally containing no phosphonomethylglycine) or PHI-F medium at pH 5.6 which comprises 4.3 g/l of MS salts (GIBCO-BRL), 0.5 mg/ ml nicotinic acid, 0.5 mg/ ml  
30 pyridoxine. HCl, 0.1 mg/ ml thiamine. HCL, 100 mg/ l myo-inositol, 2 mg/ l glycine, 40 g/ l sucrose and 1.5 g/ l gelrite.

Alternatively, selected calli are transferred directly to LSZ regeneration medium adjusted to pH 5.8 with KOH and comprising LS major and minor inorganic salts (Linsmaier and Skoog, 1965, *Physiol. Plant* 18, 100-127), 0.5 mg/ ml nicotinic acid, 0.5 mg/ ml pyridoxine. HCl, 1.0 mg/ ml thiamine. HCL, 700 mg/ l L-proline, 100 mg/ l myo-inositol, 5  
5 mg/ ml of zeatin, 20 g/ l sucrose, 0.5 g/ l MES, 250 mg/ l cefotaxime, 8 g/ l purified agar (Sigma A-7049) and, optionally, between 0.02 mM and 1 mM of tissue culture grade N-(Phosphonomethyl)-glycine (Sigma P-9556) is used. After a period of incubation in the dark plates are subject to illumination (continuous or light/day as above) and plantlets regenerated.

Small plantlets are transferred to individual glass tubes containing either PHI-F  
10 medium or half strength LSF medium at pH 5.8 comprising LS major salts (Linsmaier and Skoog, 1965, *Physiol. Plant* 18, 100-127) at half strength, LS minor salts, 0.5 mg/ ml nicotinic acid, 0.5 mg/ ml pyridoxine. HCl, 1.0 mg/ ml thiamine. HCL, 100 mg/ l myo-inositol, 20 g/ l sucrose, 0.5 g/ l MES, 8 g/ l purified agar (Sigma A-7049). and grown on for about another week. Plantlets are then transferred to pots of soil, hardened off in a growth  
15 chamber (85% relative humidity, 600 ppm CO<sub>2</sub> and 250 mE m<sup>-2</sup> s<sup>-1</sup>) and grown to maturity in a soil mixture in a greenhouse.

The first (To) generation of plants obtained as above are self fertilised to obtain second generation (T1) seeds. Alternatively (and preferably) the first generation of plants are reciprocally crossed with another non-transgenic corn inbred line in order to obtain second  
20 generation seeds. The progeny of these crosses (T1) are then expected to segregate 1:1 for the herbicide resistance trait. T1 seeds are sown, grown up in the glass house or field and the level of resistance, inheritance of resistance and segregation of resistance to the herbicide glyphosate through this and subsequent generations assessed by the observation of differential plant survival, fertility, and symptoms of necrosis in tissue following spray  
25 treatment of with glyphosate (suitably formulated and, optionally, as a salt) at a range of rates between 25 and 2000 g/ ha and at a range of growth stages between and including V2 and V8 (or, alternatively, at 7-21 days post germination). These assessments are made relative to susceptible segregants and relative to similar, untransformed lines of corn which do not comprise genes of the present or similar inventions capable of conferring resistance to  
30 glyphosate. Transgenic lines which exhibit resistance to glyphosate are selected and again selfed or backcrossed to a non-transgenic inbred.

At all stages in the above process tissue samples of transformed callus, plantlets, T0 and T1 plant material are optionally taken and analysed by 1) Southern and PCR in order to indicate the presence, copy number and integrity of transgenes, 2) Northern (or similar) analysis in order to measure expression of mRNA from transgenes, 3) quantitative Western analysis of SDS gels in order to measure expression levels of EPSPS and 4) measurement of EPSPS enzyme activity levels in the presence and absence of glyphosate in order to assess more accurately how much of the EPSPS which is expressed derives from the transgene.

Such methods of analysis are well known in the art. Suitable methods to test for the presence, integrity and expression of the transgene by PCR, for carrying out Southern analysis, for the cloning and expression of mature rice EPSPS in *E.coli*, for the purification of rice EPSPS, for the generation of polyclonal antibodies to purified rice EPSPS, for Western analysis of EPSPS levels in callus and in plant tissues and for the measurement of EPSPS activity levels in plant-derived extracts at a concentration of glyphosate which discriminates between the endogenous glyphosate-susceptible EPSPS and the glyphosate-resistant product of the EPSPS-encoding transgene are described in more detail below in Examples 17-20.

**EXAMPLE 12. Transformation of corn lines by bombardment with particles coated with DNA which includes an EPSPS expression cassette; selection and regeneration of plant cells and plants which are resistant to glyphosate**

In a further example, friable embryogenic callus derived from immature maize embryos is initiated on a solid medium and transformed biolistically. Similar to the process described in example 11, transformed callus is then selected on the basis of differential growth rate in medium containing a range of concentrations of glyphosate. Resistant callus is selected and regenerated to provide T0 plantlets which are transferred to pots, grown to maturity and self or cross fertilised in the glasshouse. The progeny seed (T1) are then grown up to provide further generations of plants which are assessed for resistance to glyphosate and analysed for transgene presence, integrity and expression as described in example 11.

**Initiation of callus from immature embryos**

Friable embryogenic Type II callus suitable for transformation is derived from immature embryos of, for example, A188 X B73 corn. Alternative inbred such as B73-derived and hybrid lines of corn can be also used including, for example, those listed in Example 11. Immature embryos of maize between 1-2 mm long are isolated aseptically from

female spikes at, typically, about 11 d after pollination using the methods indicated in example 11.

Immature embryos are plated onto, for example, onto a N6-based medium (Chu et al, 1975, Scientia Sinica, 18, 659-668) adjusted with KOH to pH 5.8 containing 1mg/ l 2,4-D, 2.9g/ l L-proline, 2 mg/ l L-glycine, 100 mg/ l of casein hydrolysate, N6 major salts, N6  
5 minor salts, N6 vitamins, 2.5 g/ l gelrite (or 2 g/ l 'Gelgro') and 20 g/ l sucrose. Alternative suitable media include, for example, a similar medium but containing MS salts (Murashige and Skoog, 1962, Physiol. Plant, 15, 473-497) in place of N6 salts. Alternatively, the medium may contain ~ 10 mg/ l dicamba in place of 2,4-D.

10 Immature embryos are incubated in the dark on the above medium at ~ 25 C in order to initiate callus. Type II callus material is selected by visual selection of fast growing friable embryogenic cells by methods known in the art and as described for example in WO 98/ 44140. For example, suitable recipient cells are selected manually by choosing preferred cells which may be at the surface of a cell cluster and further identifiable by their lack of  
15 differentiation, small size and high nucleus/ cytoplasm volume ratio. A suspension culture is initiated from tissue within the callus which appears the least differentiated, softest and most friable. Tissue with this morphology is transferred to fresh plates of media about 8- 16 d after the initial plating of the immature embryos. The tissue is then routinely subcultured every 14- 21 d by taking on ~ 10% of pieces which reach approximately a gram. At each  
20 step only material with the desired type II or type III morphology is subcultured on.

#### Preparation of cell suspension cultures

Preferably within 6 months of the above-described callus initiation, dispersed suspension cultures are initiated in liquid media containing suitable hormones such as 2,4-D and NAA optionally supplied in the form of slow-release hormone capsule treatments as  
25 described for example in examples 1 and 2 of US 5550318. Optionally, hormone levels within the cultures are maintained by occasional spiking with fresh hormone supplement. Suspension cultures are initiated, for example, by adding approximately 0.5 g of callus tissue to a 100 ml flask containing 10 ml of suspension culture medium. Every 7d, the culture is further subcultured by transferring, by use of a sterile wide-ended pipette, 1 ml of settled  
30 cells and 4 ml of conditioned medium to a fresh flask containing fresh medium. Large aggregates of cells unable to pass through the pipette tip are excluded at each subculturing step. Optionally, suspension cultures are passed through a suitable sieve (e.g. ~ 0.5-1 mm

mesh) at each subculturing step. After 6- 12 weeks the culture becomes dispersed. Suitable cell suspension culture media include for example, a medium adjusted to pH 6.0 containing Murashige and Skoog (1962) major and minor salts (optionally modified to contain a reduced level, 1.55 g/ l, of ammonium nitrate), 30 g/ l sucrose, 0.25 mg/ l thiamine, 10 mg/ l dicamba, 25 mM L-proline, 200 mg/ l casein hydrolysate, 100 mg/ l myo-inositol, 500 mg/ l potassium sulphate and 400 mg/ l potassium hydrogen phosphate. Alternatively, in place of dicamba, cell suspension medium contains 2,4-D and/or NAA.

#### Cryopreservation of cell suspension cultures

Optionally, suspension cultures obtained as described above, are cryopreserved using cryoprotectants and methods described for example in example 2 of US 5550318.

Cryopreservation entails adding cryoprotectant at ice temperature to pre-cooled cells, also at ice temperature, in a stepwise manner over a period of one to two hours. The mixture is maintained at ice temperature and the eventual volume of cryoprotectant is equal to the volume of cell suspension. The final concentrations of cryoprotectants are, for example, 10% dimethylsulfoxide, 10% polyethylene glycol (6000 Mw), 0.23 M L-proline and 0.23 M glucose. After a 30 min period of equilibration at ice temperature the mixture is divided into ~ 0.5 ml aliquots, transferred to 2 ml microcentrifuge tubes, and cooled slowly at a rate of 0.5 C/ min down to a temperature of -8 C. Following a period for ice nucleation, the sample is further cooled slowly down to -35 C and then plunged into liquid nitrogen. When required for use, frozen samples are thawed by first bathing them in their containers in water at ~ 40 C for 2 min and then allowing them to slowly thaw completely. The mixture of cells and cryoprotectants is then pipetted onto a filter laid over a layer of BMS 'feeder' cells at 25 C. Once the thawed tissue begins to grow it is transferred back to fresh solid culture medium and, once established (within 1 to 2 weeks) is further transferred into cell suspension culture medium. Once growth in liquid suspension culture is re-established the cells are used for transformation.

#### Particle-mediated transformation

Plasmid pIGPD9-derived DNA (Figure 12) containing XmaI EPSPS expression cassettes (i.e. pZEN6i, ZEN10i, etc.) is purified, bulked up (e.g by anion exchange chromatographic or CsCl<sub>2</sub> gradient densitometric isolation of plasmid DNA from cells of a suitable HisB-, Rec A- host strain of *E.coli* (e.g. DH5 $\alpha$ : hisB-) after growth to stationary phase in a minimal 5xA medium (K<sub>2</sub>HPO<sub>4</sub> 52.5g, KH<sub>2</sub>PO<sub>4</sub> 22.5g, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 5g and sodium



citrate.2H<sub>2</sub>O 2.5g per litre) and provided as a concentrated solution (preferably ~ 1 mg/ ml) in sterile water. DNA is provided as a circular plasmid DNA or, alternatively is restricted with Xma I to provide a linear EPSPS-expression cassette-containing fragment and used following purification by agarose gel electrophoresis and electroelution.

5           Suitable apparatus for bombardment is, for example, the Biorad PDS1000 Helium gun. The dish is placed 5-6 cm below the stopping screen used to stop the Kapton macroprojectile. The DNA construct is precipitated onto tungsten or gold particles with an average diameter of ~ 1.0 µm in a similar manner to that described by Klein *et al* 1987, Nature, 327, 70-73. For example, 1.25 mg of tungsten or gold particles (pre-washed in  
10 ethanol at 65 C for 12 h) are mixed, in successive order, with ~ 20-30 mg of DNA, 1.1 M CaCl<sub>2</sub> and 8.7 mM spermidine to a final volume of ~ 0.6 ml. The mixture is vortexed for 10 min at 0-4 C, subject to low speed centrifugation (~ 500g) for 5 min and the bulk of the supernatant decanted off to leave the tungsten particles suspended in a final volume of ~ 30 ml. 1- 10 µl aliquots are pipetted onto the macroprojectile of the particle gun.

15           Suspension cultures derived from type II and/or type III callus are maintained in culture for 3-5 months (or, alternatively, recovered from cryopreservation) , freshly subcultured and then sieved through a ~ 0.5-1 mm stainless steel mesh. Approximately 0.5 ml packed cell volume of cells recovered from the filtrate is then pipetted onto 5 cm paper filters and vacuum dried before transfer to a petri dish containing a stack of three 7 cm paper  
20 filters moistened with suspension culture medium. Each plate of suspension cells is centred onto the sample plate tray, the petri dish lid removed and bombarded twice at a vacuum of 28 inches of mercury. 0.1 or 1.0 mm screens are optionally placed about 2.5 cm below the stop plate in order to ameliorate injury to the bombarded tissue. After bombardment the plant cells are removed from the filter, resuspended back into cell suspension culture medium and  
25 cultured for 2-21 days. Alternatively, the bombarded callus is transferred, plate to plate, onto to a plate containing a similar solid medium (for example containing 8g/ l of purified agar) and similarly cultured at ~ 25 C in the dark.

#### Selection of transformants

Following transformation, cells growing unselected in liquid or solid culture are  
30 transferred to filters and overlayed onto solid medium containing a range (0.1 - 20 mM) of selecting concentrations of tissue culture grade N-(phosphonomethyl) glycine (Sigma). Suitable solid selection media include media, adjusted to pH 5.8 or 6.0 with KOH,

containing either MS or N6 salts (such as those described above for callus initiation or, with suitable addition of agar, those described above for growth of cells in liquid suspension) and N-(phosphonomethyl) glycine . Suitable selection media also include, for example, the selection media described in example 11 but, in this case, modified so as to lack antibiotics.

- 5 Transformed calli expressing the resistant EPSP synthase enzyme are selected on the basis of their growth at concentrations inhibitory to similar preparations of untransformed cells. Growing clumps are subcultured on to fresh selective medium. Preferably the concentration of N-(Phosphonomethyl)-glycine used in the selection medium is about 1 mM for the first two weeks of selection and about 3 mM thereafter. After 6-18 weeks putative resistant calli  
10 are identified and selected.

Regeneration of transformants/ Propagation and Analysis of transformed plant material

- The selected calli are regenerated into normal fertile plants according to, for example, the methods described by Duncan et al (1985, Planta, 165, 322-332) by Kamo et al (1985, Bot. Gaz. 146(3), 327-334) and/or by West et al (1993, The Plant Cell, 5, 1361-1369) and/or  
15 by Shillito et al (1989) Bio/ Technol. 7, 581-587.

- For example, plants are efficiently regenerated by transferring the embryogenic callus to Murashige and Skoog medium adjusted to pH 6.0 containing 0.25 mg/ l 2,4-D, 10 mg/ l 6-benzyl-aminopurine and, optionally, 0.02 to 1 mM N-(phosphonomethyl) glycine. After ~ 2 weeks tissue is transferred to a similar medium but lacking hormones. Optionally the  
20 hormone level is decreased step wise through more transfers and over a longer period of time up to 6-8 weeks. Shoots which develop after 2-4 weeks are transferred to MS medium containing 1% sucrose and solidified with 2g/ l Gelgro into which they then root.

Alternatively methods and media used for regeneration are as in example 11 except that the media used do not contain antibiotic.

- 25 Methods for growing plants to maturity, for the further propagation of plants through generations, for analysis of the inheritance of resistance to glyphosate and for analysis of the presence, integrity and expression of the EPSPS transgene are as described in example 11.

- EXAMPLE 13. Transformation of corn lines with DNA which includes an EPSPS expression cassette coated onto silicon carbide whiskers; selection and regeneration of**  
30 **plant cells and plants which are resistant to glyphosate**

In a further example, maize lines including, for example, hybrid lines having the genotype A188 x B73 are prepared as cell suspensions and transformed by contacting the

cells with silicon carbide whiskers coated with DNA using methods essentially as described by Frame et al (1994, Plant J. 6, 941-948). As described in the previous examples, the transformed callus so generated is selected on the basis of differential growth rate in medium containing a range of concentrations of glyphosate, regenerated into plantlets (To) which are grown to maturity and either self or cross fertilised to provide progeny seed (T1) for further breeding. Plants and plant material is assessed for resistance to glyphosate and analysed for transgene presence, integrity and expression as described in the previous examples.

#### Initiation of callus from immature embryos, preparation of cell suspension cultures

Maize cell suspensions suitable for transformation are optionally cryopreserved and provided in the same manner as described in example 2.

#### Transformation

Plasmid pIGPD9-derived DNA (Figure 12) containing XmaI EPSPS expression cassettes (e.g. pZEN7i, ZEN8I etc.) is purified, bulked up (e.g by anion exchange chromatographic or CsCl<sub>2</sub> gradient densitometric isolation of plasmid DNA from cells of a suitable HisB-, Rec A- host strain of *E.coli* (e.g. DH5 $\alpha$ : hisB-) after growth to stationary phase in a minimal 5xA medium (K<sub>2</sub>HPO<sub>4</sub> 52.5g, KH<sub>2</sub>PO<sub>4</sub> 22.5g, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 5g and sodium citrate.2H<sub>2</sub>O 2.5g per litre) and provided as a concentrated solution (preferably ~ 1 mg/ ml) in sterile water. DNA is provided as a circular plasmid DNA or, alternatively is restricted with Xma I to provide a linear EPSPS-expression cassette-containing fragment and used following purification by agarose gel electrophoresis and electroelution.

Transformation is carried out exactly as described by Frame et al 1994. Alternatively the procedure is somewhat modified as described below.

Cells grown in liquid culture in cell suspension medium one day after subculturing are allowed to settle out in a shake flask. Spent medium is decanted and drawn off and 12 ml of N6 medium at pH 6.0 (Chu et al 1975) modified to contain 6 mM L-proline, 20 g/ l sucrose, 2 mg/ l 2,4-D, 0.25 M sorbitol and 0.25M mannitol is added per 4 ml packed cell volume. The flask is returned to the shaker (rotary shaken at 125 rpm and incubated at 26-28 C) for 45 min. At the end of this period 1 ml aliquots of cell suspension are removed, using a wide bore pipette, into a series of sterile microcentrifuge tubes. After allowing the cells in each tube to settle out, 0.6 ml of the spent medium supernatant is then removed to leave most of the remaining content as settled cells. 50 mg of silicon carbide whiskers (Silar SC-9 whiskers, Advanced Composite Materials Corp., Greer, SC. USA) are suspended by

vortexing in 1 ml of the modified N6 medium described above. 40 µl of these suspended whiskers and 25 mg of the plasmid or linear DNA including the EPSPS expression cassette are then added to each tube of settled cells. The tubes are finger vortexed 2-3 times, mixomated (in a Mixomat dental amalgam mixer (Degussa, Ontario, Canada) for 1 second and then 0.3 ml of N6 medium (modified as described above) is added to each microcentrifuge tube. The suspended cells are then plated (200 µl/ plate) out onto a filter disc overlying solid N6 medium (the same as the modified N6 medium described above but lacking sorbitol, lacking mannitol and containing 30 g/l sucrose and 3 g/l of gelrite). Each plate is then wrapped with Urgopore tape (Stelrico, Brussels) and left to incubate in the dark for 1 week at 26-28 C.

#### Selection of transformants

Transformed callus is selected as described in example 12 or, alternatively, as described in Frame et al 1994 except that N-(phosphonomethyl)glycine is used, at a range of concentrations between 1 and 5 mM in place of the bialaphos specified in the Frame et al publication.

#### Regeneration of transformants/ propagation and analysis of transformed plant material

Plants are regenerated, propagated, and bred as described in example 12. Plants are analysed for resistance to glyphosate and plant material is analysed for transgene presence, integrity and expression as described in example 12

#### **TABLE 2. Expression of EPSPS transgene in regenerable callus following transformation using silicon carbide Whiskers**

The table shows EPSPS enzyme assay (+/- 100 µM glyphosate at 100 µM PEP) results based upon enzyme assays of extracts of stably transformed callus of regenerable A188 x B73 regenerable corn, transformed by Whiskers with ZEN13 DNA. Each callus line represents a single event which is assayed in duplicate. The ratio of the true (allowing for ~ 8% inhibition) tolerant enzyme activity (expressed by the transgene) to endogenous susceptible activity (>98% inhibition + glyphosate) is calculated. The mutant EPSPS is expressed relatively strongly in one particular line, 90921sw3-1, where, allowing for the reduced Vmax of the tolerant enzyme relative to the w/t (about a third) it can be estimated that the tolerant enzyme is expressed at 3-10X the normal level of endogenous EPSPS (this calculation is complicated by the fact that in this particular event the endogenous susceptible level of EPSPS activity appears unusually low). The same extracts were also analysed by

Westerns (in this case using polyclonal antibodies raised to purified *Brassica napus* EPSPS) and the amount of EPSPS quantitated on the basis of reaction with a standard curve of purified rice EPSPS. The Western data are expressed as fold increase in total EPSPS amount relative to untransformed corn callus. In good agreement with the enzyme data, the Western data indicate a high level of EPSPS expression in, for example, lines 90928sw3-1.

Event. Line#	DNA Construct	Measured activity (nmol/ min/ mg) + 100uM glyphosate (true tolerant activity = measured x 1.08)	Total Activity (nmole / min / mg) in absence of glyphosate	Ratio of (true) tolerant/ sensitive EPSPS activity	Western analysis(X- fold relative to control)
90921sl5-1	ZEN13	2.87	17.04	1:4	4
		3.14	11.84		
90921tl1-1	ZEN13	1.66	10.89	1:6	3
		2.03	15.88		
90928tl2-1	ZEN13	2.61	20.04	1:4.5	3
		4.3	15.86		
90921sw3- I	ZEN13	11	13.22	1:0.3	7
		8.88	14.96		

**EXAMPLE 14. Transformation of rice lines using an *Agrobacterium* strain containing a superbinary vector which includes an EPSPS expression cassette between the right and left borders of the T-DNA; selection and regeneration of plant cells and plants which are resistant to glyphosate**

In a further example, scutella are isolated from mature seeds of suitable lines of rice (including, for example, varieties Koshihikari, Tsukinohikari and Asanohikari) dedifferentiated and the callus thus-obtained transformed by infection with *Agrobacterium*. Following selection and regeneration, transgenic plantlets (To) are obtained which are grown to maturity and either self or cross fertilised to provide progeny seed (T1) for further breeding. Plants and plant material are assessed for resistance to glyphosate and analysed for transgene presence, integrity and expression as described in the previous examples. As an

alternative to the methods described below the methods described in example 1 of US 5591616, suitable adapted so that glyphosate rather than hygromycin is used for selection, are used.

Construction of *Agrobacterium* strain; Preparation of *Agrobacterium* suspension

5 A strain of *Agrobacterium* containing superbinary vector having the desired EPSPS expression cassette between the right and left borders is constructed (using electroporation to transform *Agrobacterium* with plasmid DNA) as described in example 11. Suspensions are prepared according to the methods described in example 11. Alternatively, the transformed strain of *Agrobacterium* is grown for 3 days on AB medium (Chilton et al, 1974, Proc. Natl. Acad. Sci. USA, 71, 3672-3676) containing appropriate antibiotic selection (e.g. 50 mg/ l spectinomycin in the case of LBA4404 (pSB1ZEN13 etc)) and looped off of the plate to form a suspension in AAM medium (Hiei et al, 1994, The Plant Journal, 6(2), 271-282) at a density of  $1-5 \times 10^9$  cells/ ml.

Rice cultivars, preparation of callus from scutella

15 Rice cultivars are, for example *Oryza sativa* L. Tsukinohikari, Asanohikari and Koshihikari.

Mature seeds are dehusked, surface sterilized by washing in 70% ethanol and then soaked for 30 minutes in 1.5% NaOCl. After rinsing in sterile water they are cultured at 30 C, in darkness for 3 weeks on 2N6 medium at pH 5.8 which contains the major salts, minor salts and vitamins of N6 medium (Chu 1978 in Proc. Symp. Plant Tissue Culture., Peking: Science Press, pp 43-50) 30 g/ l sucrose, 1 g/ l casein hydrolysate, 2 mg/ l 2,4-D and 2 g/ l gelrite. Proliferated callus derived from the seed scutella is subcultured for 3-7 days on fresh 2N6 medium. Growing callus (1-2 mm in diameter) is selected, suspended in 2N6 liquid medium (without gelrite) and cultured in flasks, in darkness on a rotary shaker at 125 rpm and at 25 C. The medium is changed every 7 days. Cells growing in log phase after 3- 4 transformations are used for transformation.

Infection, transformation and selection

Suspended rice callus cells are allowed to settle out of suspension and then resuspended in the suspension of *Agrobacterium*, left in contact for several minutes and then, again, allowed to settle out and, without rinsing, plated out onto 2N6-AS medium (2N6 medium adjusted to pH 5.2 and containing 10 g/ l D-glucose and 100  $\mu$ M acetosyringone) and incubated in the dark at 25 C for 3-5 days. Growing material is rinsed thoroughly with

250 mg/ l cefotaxime in sterile water and then transferred onto 2N6-CH medium (2N6 medium adjusted to pH 5.8 with KOH containing 250 mg/ l cefotaxime and 0.5 - 5 mM tissue culture grade N-(phosphonomethyl) glycine) or, alternatively, 2N6K-CH medium (2N6 medium modified as described by Hiei et al 1994 but, in place of hygromycin, containing 0.5  
5 - 5 mM tissue culture grade N-(phosphonomethyl) glycine) and cultured for 3 weeks in the dark at 25C. Proliferating colonies are subcultured onto a second plate of selective medium for a further period of 7-14 days.

#### Regeneration and analysis of plants

Growing colonies are plated onto a regeneration medium at pH 5.8 containing half  
10 strength N6 major salts, N6 minor salts, N6 amino acids, vitamins of AA medium (Chilton et al 1974), 1 g/ l casein hydrolysate, 20 g/ l sucrose, 0.2 mg/ l naphthaleneacetic acid, 1 mg/ l kinetin, 3 g/ l gelrite and, optionally, 0.04-0.1 mM N-(phosphonomethyl)glycine. These plates are incubated at 25 C and kept under continuous illumination (~ 2000 lux). As described in example 1 regenerated plants are eventually transferred to soil in pots and  
15 matured in a greenhouse.

Plants are propagated, and bred (for example the transgenic plants are selfed) essentially as described in example 11. Plants are analysed for resistance to glyphosate and plant material is analysed for transgene presence, integrity and expression essentially as described in example 11.

#### **EXAMPLE 15. Transformation of wheat lines with DNA which includes an EPSPS expression cassette by use of microprojectile bombardment; selection and regeneration of plant cells and plants which are resistant to glyphosate**

In a further example, immature embryos are isolated from suitable lines of wheat (including, for example, spring wheat cv BobWhite, and Jaggar) incubated on hormone( 2,4-  
25 D) -containing medium for 2 days and transformed by bombardment with DNA-coated particles. Following a period for recovery and continued growth of callus, callusing embryos are subcultured through a series of media containing a fixed level of glyphosate and (serially diluted) decreasing levels of 2,4-D such that somatic embryogenesis is induced. The selected material is regenerated to form shoots on a medium also containing glyphosate,  
30 transferred to rooting medium and, as in the previous maize-related examples, regenerated into plantlets (To) which are grown to maturity and either self or cross fertilised to provide progeny seed (T1) for further breeding. Plants and plant material are assessed for resistance

to glyphosate and analysed for transgene presence, integrity and expression as described in the previous examples. As an alternative to the methods described below the methods described in example 1 of US 5631152 are used.

#### Preparation of immature embryos

5 Wheat plant lines (for example spring wheat *Triticum aestivum* cv BobWhite) are grown to maturity in the greenhouse and caryopses isolated at 11 - 15 postanthesis. Caryopses are surface sterilised by treatment for 15 minutes in 5%NaOCl and then washed repeatedly in sterile water. Immature embryos are aseptically isolated onto 3cm squares of nylon netting (mesh size 1.5 mm) overlying A2 medium. A2 medium adjusted to pH 5.8 is  
10 4.32 g/l Murashige and Skoog salts, 20 g/l sucrose, 0.5 g/l L-glutamine, 2 mg/l 2,4-D, 100 mg/l casein hydrolysate, 2 mg/l glycine, 100 mg/l myo-inositol, 0.5 mg/l nicotinic acid, 0.1 mg/l thiamine.HCl and 2.5 g/l gelrite. Embryos are arranged into a solid 2.5 cm disc, comprising approx. 50 in number. Plates are sealed with leukopore tape and incubated at 25°C in the dark for 2 days. Four hours prior to bombardment embryos are transferred onto  
15 plates containing fresh A2 medium supplemented with 36.44 g/l D-sorbitol and 36.44 g/l D-mannitol. The embryos are transferred from plate to plate by means of the nylon net upon which they sit. The embryos sit on this increased osmotic strength medium for 4 h at 25°C in the dark before being bombarded.

#### Particle-mediated transformation

20 Plasmid pIGPD9-derived DNA (Figure 12) containing XmaI EPSPS expression cassettes (i.e. pZEN6i, ZEN10I etc.) is purified, bulked up (e.g by anion exchange chromatographic or CsCl<sub>2</sub> gradient densitometric isolation of plasmid DNA from cells of a suitable HisB-, Rec A- host strain of *E.coli* (e.g. DH5α: hisB-) after growth to stationary phase in minimal 5xA medium (K<sub>2</sub>HPO<sub>4</sub> 52.5g, KH<sub>2</sub>PO<sub>4</sub> 22.5g, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> 5g and sodium  
25 citrate.2H<sub>2</sub>O 2.5g per litre) and provided as a concentrated solution (preferably ~ 1 mg/ml) in sterile water. DNA is provided as a circular plasmid DNA or, alternatively is restricted with Xma I to provide a linear EPSPS-expression cassette-containing fragment following purification by agarose gel electrophoresis and electroelution.

Particles are prepared and coated with DNA in a similar manner to that described by  
30 Klein *et al* 1987, Nature, 327, 70-73. Preparation of DNA-coated particles and operation of the particle gun is as described in example 12. Alternatively, the details are as follows. For example, 60 mg of gold or tungsten particles (~ 1.0 µm) in a microcentrifuge tube are washed



repeatedly in HPLC-grade ethanol and then, repeatedly, in sterile water. The particles are resuspended in 1 ml of sterile water and dispensed into 50 µl aliquots in microcentrifuge tubes. Gold particles are stored at 4 C, tungsten particles at - 20 C. 3 mg of DNA are added to each aliquot of (defrosted) particles and the tubes are vortexed at top speed. Whilst  
5 maintaining near continuous vortexing, 50 µl of 2.5M CaCl<sub>2</sub> and 20 µl of 0.1M spermidine is added. After 10 minutes of further vortexing, samples are centrifuged for 5 seconds in an eppendorf microcentrifuge, the supernatant is drawn off and the particles washed in successive additions of HPLC-grade ethanol. The particles are thoroughly resuspended in 60 µl of ethanol and then dispensed in 10 µl aliquots onto the surface of each kapton membrane  
10 macrocarrier to be used in the PDS1000 particle gun.

Components of the PDS1000 particle gun are surface sterilised by immersion in 70% ethanol and air-drying. Target plates prepared, as described above, with ~ 50 embryos arranged into an ~ 2.5 cm disc are placed 6 cm from the stopping screen. 1100 psi rupture discs are then used for bombardment. Each plate is bombarded once or twice.

15 Bombarded plates are sealed with pore tape and maintained at 25 C in the dark for ~ 16h. Embryos dislodged from the surface of the medium by the helium shock wave are recovered and also incubated overnight on fresh plates of the same mannitol and sorbitol-supplemented A2 medium. The bombarded embryos are then transferred to fresh plates of A2 medium and incubated for 1 week at 25 C in the dark prior to selection.

#### 20 Selection and Regeneration of Transformants

After this recovery period callusing embryos are removed from the nets and transferred to A2 2P medium (A2 medium, adjusted to pH 5.8 containing 2 mM N-(phosphonomethyl)glycine), at a density of 20 explants / plate. After one week on A2 2P medium, calli are removed to A1 2P medium (A2 medium containing only 1.0 mg / l 2,4-D  
25 and 2 mM N-(phosphonomethyl)glycine) for 2 weeks and thence to A 0.5 2P medium (A2 medium containing only 0.5 mg / l 2,4-D and 2 mM N-(phosphonomethyl)glycine) for a further two weeks. Optionally, the 2 week incubation periods are reduced to 1 week and/or the middle step of incubation on A1 2P medium is omitted. Optionally, the selecting concentration of N-phosphonomethylglycine is between 0.5 and 10 mM although 2 mM is  
30 preferred. The overall time for this period of callus induction with descending levels of 2,4-D in the medium is 2- 10 weeks, preferably 3-6 weeks and most preferably ~ 4 weeks.

To encourage maximum shoot growth and to discourage root development the calli are then transferred to Z medium. Z medium is A2 medium but containing 10 mg/ l zeatin in place of 2,4-D and also containing 0.1 mM N-(phosphonomethyl)glycine. Optionally, N-(phosphonomethyl)glycine is in the range 0.04 - 0.25 mM. Regenerating calli are  
5 maintained on this medium for a period of 3 weeks before subculture, at which point well developed shoots are excised. As only one event is likely to be produced on a single callus (which represents a single embryo), the entire callus is removed to a fresh plate and maintained with the excised shoot(s) to ensure multiple clones arising from the same callus do not get counted as separate events. Calli with only partially developed shoots or without  
10 regenerating sectors are returned to Z medium for a further 3 weeks. At the end of this period non regenerating calli are discarded.

Shoots are maintained on Z medium until 4 or more well-developed leaves (extending to ~ 2 cm in length) have formed. The regenerating plant material is then carefully transferred to plastic tubs containing 0.5MS medium. 0.5 MS medium at pH 5.8 is 2.16 g/ l  
15 of Murashige and Skoog salts, 15 g/ l sucrose, 2.5 g activated charcoal, 2.5 g/ l gelrite, 1 mg/ l glycine, 50 mg/ l myo-inositol, 0.25 mg/ l nicotinic acid, 0.25 mg/ l pyridoxine.HCL, 0.05 mg/ l thiamine.HCl and 0.1 mM N-(phosphonomethyl)glycine (optionally 0.0-0.25 mM).

Once plants have rooted they may be potted into soil and weaned, or removed to individual glass boiling tubes containing 0.5MS (with no N-(phosphonomethyl)glycine) and  
20 2.5 g / l charcoal. It is preferred to have charcoal present in the rooting medium to adsorb any remaining PGRs or selection chemical transferred with the plantlet, and to create a dark rooting environment thereby avoiding physiologically aberrant green roots.

Callus induction and the first week of regeneration occurs at 25°C in the dark. The second week of regeneration occurs at low light at 25°C, then subsequent weeks at approx.  
25 2500 lux on a 16 hour photo period.

#### Propagation, breeding and analysis of transformed plant material

Methods of producing T1 and further progeny are all well known in the art and essentially as described in the previous examples. Methods for analysis of the inheritance of resistance to glyphosate and the presence, integrity and expression of transgenes are as in the  
30 previous examples

**EXAMPLE 16. Transformation of wheat lines with DNA which includes an EPSPS expression cassette by electroporation of protoplasts; selection and regeneration of plant cells and plants which are resistant to glyphosate**

In a further example, plasmid or linear DNA comprising an EPSPS expression cassette and identical to that used in examples 12, 13 and 15 is used for direct transformation of protoplasts of a line of wheat capable of regeneration into fertile plants (*cf* US 5231019). Isolated protoplasts of wheat, preferably from leaf tissue or cells in culture (*cf* Gamborg, O.L. and Wetter, L.R., Plant Tissue Culture Methods, 1975, 11-21) are prepared at ~ ca  $2 \times 10^6$  protoplasts/ml in 0.4M mannitol at pH 5.8. To this suspension are added first, 0.5 ml of 40% w/v polyethylene glycol (PEG) of molecular weight 6000 in modified F medium (Nature (1982), 296, 72-74) at pH 5.8 and, second, 65 ml of water containing 15 mg of the desired plasmid or linear DNA and 50 mg of calf thymus DNA. The mixture is incubated together for 30 min at 26 C, occasionally agitated and subsequently diluted into F medium (Nature (1982), 296, 72-74). The protoplast are isolated by low-speed centrifugation, taken up in 4 ml of CC culture medium (Potrykus, Harms, Lorz, (1979) Theor. Appl. Genet., 54, 209-214) and incubated in the dark at 24 C.

Alternatively, and in addition to treatment with PEG, transformation of cereal protoplasts is carried out using further steps of heat shock and/or electroporation (Neumann, E. et al (1982), the EMBO J. , 7, 841-845). Thus, for example, wheat protoplasts are incubated in an aqueous solution of DNA and mannitol, heated to 45 C for 5 min and then cooled to 0 C over a period of 10 seconds. Then polyethylene glycol is added (Mr 3K -8K) until the final concentration is ~ 8% w/v. After gentle but thorough mixing treatment is carried out in an electroporator. The chamber of a Dialog 'Porator' (Dialog, Dusseldorf, Germany) is sterilised by washing with 70% ethanol and then drying in sterile air. Suspensions of protoplasts (~ ca  $2 \times 10^6$  protoplasts/ml in 0.4M mannitol + the DNA ) are adjusted with manganese chloride to a measured electrical resistance of ~ 1.4 k ohm. Samples of volume ~ 0.4 ml are subjected, at 10 second intervals, to three pulses of applied voltages of between 1000 and 2000 V. The, thus transformed protoplasts are then collected and diluted back out into CC culture medium.

Those skilled in the art will recognise that many permutations and variations of these transformation procedures are possible and that, for example, transformation may also be

improved by raising the pH to 9.5 and/or increasing the concentration of calcium ions in the solution within which transformation is carried out.

After 3-14 days aliquots of the developing cell cultures are transferred to medium containing alternative selecting concentrations of tissue-culture grade N-(phosphonomethyl) glycine (Sigma) between 1 and 5 mM (preferably 2 mM). Resistant cell colonies so identified (exhibiting growth on concentrations of glyphosate at least 2 fold greater than tolerated by untransformed controls) are transferred to fresh agar medium also containing a range of selecting concentrations of glyphosate and, as described, in example 15, subcultured between plates containing successively declining concentrations of 2,4-D. Growing resistant colonies May be analysed (by PCR etc) for the presence of the recombinant DNA. It may or may not be possible to effect much selection at the callus step. In any case all growing calli will be taken forward.

Growing calli are then transferred to shoot regeneration medium containing zeatin and N-(phosphonomethyl)glycine and thence to rooting medium exactly as described in example 15. Fertile transgenic plants expressing glyphosate-resistant EPSP synthase are then regenerated, selected and tested as known in the art and as described in example 15 and using the analytical methods described in example 11.

**EXAMPLE. 17. Method for assaying EPSPS activity and determination of kinetic constants. Method for assaying EPSPS activities in crude plant materials and discriminating the proportion of the total which is resistant to glyphosate.**  
EPSPS Enzyme assay

Assays are carried out generally according to the radiochemical method of Padgett et al 1987 (Archives of Biochemistry and Biophysics, 258(2) 564-573) with K<sup>+</sup> ions as the major species of cationic counterion. Assays in a total volume of 50µl, in 50mM Hepes(KOH) pH 7.0 at 25°C, contain purified enzyme or plant extract (see below) diluted appropriately in Hepes pH 7.0 containing 10% glycerol, and 5mM DTT, <sup>14</sup>C PEP either as variable substrate( for kinetic determinations) or fixed at 100 or 250 µM and shikimate 3 Phosphate (K<sup>+</sup> salt) at 2 or 0.75 mM as indicated. Optionally, for assays of crude plant extracts, assays also contain 5 mM KF and/or 0.1 mM ammonium molybdate. Assays are started with the addition of <sup>14</sup>C phosphoenolpyruvate (cyclohexylammonium+ salt) and stopped after 2-10 minutes (2 minutes is preferable) with the addition of 50µl of a solution of 1 part 1M acetic acid and 9 parts ethanol. After stopping, 20µl is loaded onto a synchropak AX100 ( 25cm x 4.6mm )

column and chromatographed using isocratic elution with a 0.28M potassium phosphate pH 6.5 mobile phase flowing at 0.5 ml/min over 35 minutes. Under these conditions the retention times for PEP and EPSP are ~ 19 and 25 minutes respectively. A CP 525TR scintillation counter is connected to the end of the AX 100 column. It is fitted with a 0.5ml flow cell, and the flow rate of scintillant ( Ultima Flo AP ) is set at 1ml/min. Relative peak areas of PEP and EPSP are integrated to determine the percentage conversion of labelled PEP to EPSP. Apparent  $K_m$  and  $V_{max}$  values are determined by least squares fit to a hyperbola with simple weighting using the Grafit 3.09b from Erithacus Software Ltd.  $K_m$  values are generally ascertained using 8-9 concentrations of variable substrate ranging from  $K_m / 2$  -  $10 K_m$  and triplicate points. Except where specifically noted, data points are only included in the analysis where there is < 30% conversion of substrate to EPSP.

Shikimate-3-Pi (S3P) is prepared as follows, To 7mls of 0.3M TAPS pH 8.5 containing 0.05M Shikimate, 0.0665M ATP ( Na salt ), 10mM KF, 5mM DTT, and 0.05M  $MgCl_2 \cdot 6H_2O$ , 75 $\mu$ l of a 77 unit (  $\mu$ mol min<sup>-1</sup> ) ml<sup>-1</sup> solution of shikimate kinase is added. After 24hrs at room temperature, the reaction is stopped by brief heating to 95°C. The reaction solution is diluted 50 fold in 0.01M Tris HCl pH 9, and chromatographed by anion exchange on Dowex 1 X 8 - 400, using a 0 - 0.34M  $LiCl_2$  gradient. The S3P fractions are combined, freeze dried, and then redissolved in 7mls distilled  $H_2O$ . 28mls of 0.1M  $Ba(CH_3COOH)_2$  and 189mls of absolute ethanol are then added. This solution is left to stir overnight at 4°C. The resulting precipitate of tri-Barium S3P is collected and washed in 30mls of 67% ethanol. The washed precipitate is then dissolved in ~ 30mls distilled  $H_2O$ . By adding either  $K_2SO_4$  the  $K^+$  or  $TMA^+$  salt of S3P is produced as required. Great care is taken to add a minimal excess of sulphate. The  $BaSO_4$  precipitate is removed and the supernatant containing the required salt of S3P freeze dried. Each salt is weighed and analysed by proton NMR. S3P preparations so-prepared are > 90% pure according to proton NMR and (according to their weights and integration of  $^{31}P$  NMR) contain only low residues of potassium sulphate.

#### Preparation of extracts of plant material suitable for EPSPS assay

Callus or plantlet material (0.5 - 1.0 g) is ground to a fine frozen powder in a liquid nitrogen-chilled mortar and pestle. This powder is taken up in an equal volume of a suitable chilled extraction buffer (for example, 50 mM Hepes/ KOH buffer at pH 7.5 containing 1 mM EDTA, 3 mM DTT, 1.7 mM 'pefabloc' (serine protease inhibitor), 1.5 mM leupeptin,

1.5 mM pepstatin A, 10% v/v glycerol and 1% polyvinylpyrrolidone), resuspended, mixed and centrifuged in a chilled centrifuge to bring down debris. The supernatant is exchanged down a chilled PD10 column of Sephadex G25 into 25 mM Hepes/ KOH buffer at pH 7.5 containing 1 mM EDTA, 3 mM DTT and 10% v/v glycerol. Protein is estimated by the  
 5 Bradford method standardised using bovine serum albumen. A portion of the extract is frozen in liquid nitrogen; a portion is assayed immediately.

EPSPS assays of plant extracts are standardly carried out, as described above, with 0.1 mM  $^{14}\text{C}$ -PEP and 0.75 mM shikimate-3-Pi either in the absence or the presence of 0.1 mM N-(phosphonomethyl)glycine. Under these assay conditions, the resistant form of  
 10 EPSPS (see below) is estimated to be inhibited by < 8.5% whilst the sensitive w/t form is essentially fully inhibited (> 98%). Thus, the level of activity observed in the presence of glyphosate (A) is taken to represent ~ 92% of the level of resistant enzyme derived from expression of the transgene whilst the level of susceptible w/t EPSPS is taken to be the total level of EPSPS activity observed in the absence of glyphosate minus the value of A x ~ 1.08.  
 15 Because the Vmax of the mutant enzyme is estimated to be only about a third of the Vmax of the w/t enzyme (and because the Km values for PEP of both w/t and mutant forms are estimated to be about 20  $\mu\text{M}$  or less), the level of expression of the mutant enzyme polypeptide relative to the level of expression of the endogenous w/t EPSPS is taken to be about three fold higher than the ratio calculated on the basis of the ratio of their relative  
 20 observed activities. The total level of EPSPS polypeptide expression (mutant + w/t) is also estimated by using Westerns (see below).

**EXAMPLE. 18. Cloning and expression of w/t and mutated cDNA encoding mature rice EPSPS in *E.coli*. Purification and characterisation of w/t and mutant rice EPSPS**  
**Expression, purification and characterisation of w/t mature rice EPSPS**

25 Rice EPSPS cDNA is amplified using RT-PCR from RNA isolated from rice variety Koshihikari using Superscript RT from BRL according to the recommendation supplied by the manufacturer. PCR is performed using Pfu turbo polymerase from Stratagene according to the methods supplied by the manufacturer. The oligonucleotides below are used in the amplification reaction and the reverse transcription steps.

30 SEQ. ID. NO. 31

Rice 3' oligo 5' GCGCTCGAGTCAGTTCCTGACGAAAGTGCTTAGAACGTCG 3'

SEQ. ID. NO. 32

Rice 5' oligo 5' GCGCATATGAAGGCGGAGGAGATCGTGC 3'

The PCR product is cloned into pCRBlunt II using Invitrogens Zero Blunt TOPO kit. The sequence of the insert is confirmed by sequencing and it is verified that the predicted open reading frame corresponds to that of the predicted mature chloroplastic rice EPSPS protein with the exception of the presence of an initiating Met. The cloned and verified rice epsps sequence is excised using Nde I and Xho I and the purified fragment is cloned into pET24a (Novagen) digested similarly. The recombinant clones are introduced into BL21 (DE3) a codon-optimised RP strain of *E.coli* supplied by Stratagene. The EPSPS protein is expressed in this strain following addition of 1 mM IPTG to the fermenter medium (LB supplemented with 100µg/ml Kanamycin). The recombinant protein of the correct predicted mass is identified i) on the basis of Coomassie staining of SDS gels of cell extracts and side by side comparison with Coomassie-stained gels of extracts of similar *E.coli* cells transformed with an empty pET24a vector and ii) by western analysis using a polyclonal antibody raised to previously-purified plant EPSPS protein. The mature rice EPSPS protein is purified at ~ 4 C as follows. 25 g wet weight of cells are washed in 50 ml of 0.1M Hepes/ KOH buffer at pH 7.5 containing 5 mM DTT, 2 mM EDTA and 20% v/v glycerol. Following low-speed centrifugation, the cell pellet is resuspended in 50 ml of the same buffer but also containing 2 mM of 'Pefabloc' a serine protease inhibitor. Cells are evenly suspended using a glass homogenizer and then disrupted at 10000 psi using a Constant Systems (Budbrooke Rd, Warwick, U.K.) Basic Z cell disrupter. The crude extract is centrifuged at ~ 30,000 g for 1 h and the pellet discarded. Protamine sulphate (salmine) is added to a final concentration of 0.2% , mixed and the solution left to stand for 30 min. Precipitated material is removed by centrifugation for 30 min at ~ 30,000 g. Aristar grade ammonium sulfate is added to a final concentration of 40% of saturation, stirred for 30 min and then centrifuged at ~ 27,000 g for 30 min. The pellet is resuspended in ~ 10 ml of the same buffer as used for cell disruption, further ammonium sulfate is added to bring the solution to ~ 70% of saturation, the solution is stirred for 30 min and centrifuged again to yield a pellet which is resuspended in ~ 15 ml of S200 buffer (10 mM Hepes/ KOH (pH 7.8) containing 1 mM DTT, 1 mM EDTA and 20% v/v glycerol). This is filtered (0.45 micron) loaded and chromatographed down a K26/ 60 column containing Superdex 200 equilibrated with S200 buffer. EPSPS-containing fractions detected on the basis of EPSPS enzyme activity are combined and loaded onto an xk16 column containing 20 ml of HP Q-Sepharose equilibrated with S200 buffer. The column is

washed with S200 buffer and then EPSPS eluted within a linear gradient developed from 0.0M to 0.2M KCl in the same buffer. EPSPS elutes within a single peak corresponding to a salt concentration at or below 0.1 M. EPSPS-containing fractions detected on the basis of EPSPS enzyme activity are combined and loaded onto a HiLoad xk26/60 column of Superdex 75 equilibrated with Superdex 75 buffer (25 mM Hepes/ KOH (pH 7.5) containing 2 mM DTT, 1 mM EDTA and 10% v/v glycerol). EPSPS elutes later from the column than might be expected on the basis of the molecular weight of the presumed dimer. This may be due to interaction of the protein with the gel matrix at low ionic strength. EPSPS-containing fractions identified on the basis of enzyme activity are combined and loaded onto a 1 ml column of MonoQ equilibrated with the same, Superdex 75 buffer. The column is washed with starting buffer and EPSPS eluted as a single peak over the course of a 15 ml linear gradient developed between 0.0 and 0.2M KCl. EPSPS is obtained near (>90% pure) at this stage in the purification. Optionally, EPSPS is further purified by exchange into Superdex 75 buffer containing 1.0 M (Aristar) ammonium sulphate and loading onto a 10 ml column of phenyl sepharose equilibrated in the same buffer. EPSPS is eluted as a single peak early during the course of a linear gradient of declining ammonium sulphate developed between 1.0 and 0.0 M ammonium sulphate.

Cloning, expression, purification and characterisation of glyphosate-resistant mutant rice EPSPS

The rice EPSPS cDNA in pCRBlunt is used as a template for two further PCR using the following primer pairs designed to introduce specific changes

SEQ ID NO 33

rice 5' oligo 5' GCGCATATGAAGGCGGAGGAGATCGTGC 3'

SEQ ID NO 34

Rice mutant reverse to RV

5' GCAGTCACGGCTGCTGTCATGATCGCATTGCAATTCCAGCGTTCC 3'

SEQ ID NO 35

Rice 3' oligo 5' GCGCTCGAGTCAGTTCCTGACGAAAGTGCTTAGAACGTCG 3'

SEQ ID NO 36

Rice mutant forward to sal 5' GGAACGCTGGAATTGCAATGCGATCATTGACAGCAGCCGTGACTGC 3'

The resultant products are gel purified and placed into a tube in equi-molar concentrations to serve as a template for another round of PCRs with the rice 5' and 3' oligos. The resultant



products are cloned into pCRBlunt II using Invitrogens Zero Blunt TOPO kit. It is confirmed that the DNA sequence of the insert and its predicted open reading frame correspond to that of the predicted mature chloroplastic rice EPSPS protein (with the exception of the presence of an initiating Met) and also that the desired changes (the specific mutation of T to I and P to S at specific positions in the EPSPS sequence) are encoded. The thus cloned and verified rice epsps sequence is excised using Nde I and Xho I and the purified fragment cloned into pET24a (Novagen) digested similarly. The recombinant clones are introduced into BL21 (DE3), a codon optimised RP strain of *E.coli* supplied by Strategene. The EPSPS protein is expressed in this strain following addition of 1 mM IPTG to the fermenter medium (LB supplemented with 100ug/ml Kanamycin). The recombinant protein of the correct predicted mass is identified i) on the basis of Coomassie staining of SDS gels of cell extracts and side by side comparison with Coomassie-stained gels of extracts of similar *E.coli* cells transformed with an empty pET24a vector and ii) by western analysis using a polyclonal antibody raised to previously-purified plant EPSPS protein. This mutant form of rice EPSPS is purified and characterised in a similar manner to the method described above for w/t rice EPSPS.

The so-obtained mutant form of rice EPSPS, assayed as described above in the presence of 2 mM shikimate-3-Pi, has an apparent  $V_{max}$  of  $\sim 10 \mu\text{mol}/\text{min}/\text{mg}$  and a  $K_m$  for PEP of  $22 \mu\text{M}$ . At  $40 \mu\text{M}$  PEP, the  $IC_{50}$  value for the potassium salt of glyphosate is  $\sim 0.6 \text{ mM}$ . The estimated  $K_i$  value for potassium glyphosate of the mutant EPSPS is  $\sim 0.2 \text{ mM}$ .

**EXAMPLE. 19. Preparation of antibodies to purified rice EPSPS and methods for Western analysis**

Standard methods for generation of polyclonal antisera in rabbits are used. Rabbits are young female New Zealand Whites. Immunisation courses consist of 4 doses, each  $\sim 100 \text{ mg}$ , administered at monthly intervals. Each dose in phosphate buffered saline is administered as an emulsion with Freund's Complete adjuvant (dose 1) or Incomplete adjuvant (doses 2-4) and is injected at four sub-cutaneous sites. A pre-bleed is taken before dose 1 is administered. A test bleed is taken 14 days after the second dose to confirm the immune response. A term bleed is taken 14 days after the fourth dose and used for experimentation. A fifth and final dose is given when at least 6 weeks has elapsed since the fourth dose, and the final bleed (also used for experimentation) is taken 14 days later.

Antibodies are raised to both (i) purified native mature w/t rice EPSPS (example 8) and also (ii) to SDS-denatured rice EPSPS polypeptide which is eluted from a band cut out of a 12% SDS gel (the correct position of the protein being accurately marked by side by side Coomassie staining of the band).

5           12% polyacrylamide gels are used for SDS gel electrophoresis and Western blotting. Electrophoresis is performed at a constant current of 100V for 90 minutes. Gels are blotted against nitrocellulose sheets overnight at 4 C at a constant 30V. Sheets are blocked in 5% Marvel phosphate buffered saline containing 0.1% Tween (PBS-tween) for 1 or 2 hours, washed three times in PBS-tween and incubated in rice EPSPS-Rb1 primary antibody at ~  
10   1.3 mg IgG/ml (normally equivalent to a 1:4000 to 1: 20,000 dilution of term bleed). These antibody dilutions are applied in PBS (phosphate-buffered saline) containing 1% Marvel and 0.05% Tween 20 for 2 hours. Secondary antibody, Goat anti Rabbit HRP (Sigma A6154) is applied at 1:10,000 or 1:20,000 in PBS containing 0.05% Tween and 1% Marvel. Incubation with secondary antibody is continued for 1 hour, the blott is washed three times in PBS  
15 (0.05% tween), ECL substrate is applied for usually 1 minute and film exposed for 10-60 seconds. Negative control blots are blots with (1) preimmune serum at a dilution calculated to yield the same [IgG] as in test immune sera (IgG is routinely purified from an aliquot of each serum and quantitated so that these dilutions can be calculated directly) and also (2) immune serum raised against Freund's adjuvant alone. IgG concentration in control immune  
20 sera is adjusted so that controls are at an appropriate concentration of IgG. In order to make this calculation, IgG is purified from crude antiserum by filtration through a 0.45µm syringe filter and passed down a 1ml HiTrap protein G column (Pharmacia cat no:17-0404-01). The bound IgG is eluted from the column with 0.1M glycine HCl pH 2.9 and dialysed vs PBS overnight. The IgG concentration is estimated by UV determination. (a 1 cm pathlength of a  
25 1 mg ml<sup>-1</sup> solution of pure IgG has an absorbance at 280nm wavelength of 1.35). From the IgG yield a calculation can be made of IgG concentration in the crude antiserum, and correct dilutions in western blots calculated accordingly.

Plant tissue samples are prepared for example as described in example 17. Alternatively, for Western analysis, a much simpler procedure is used. 100-200 mg of plant  
30 tissue to be analysed is rapidly homogenized (for example using an ultra turrax, bead beater or glass homogenizer) in an equal volume of buffer (similar to in example 7), centrifuged for 5 minutes in a chilled eppendorf microcentrifuge and the supernatant a) a small aliquot is

analysed for protein using the Bradford method and b) for the most part mixed 1:1 with Laemli SDS 'sample buffer', heated and then stored ready for loading onto gels. Typically SDS slab gels are loaded with 10 protein samples in 10 wells. Typically these are 1 to 10 µg of crude extracts of plant material for analysis alongside a standard curve of between 0 and 20 ng of pure rice EPSPS. In some cases Westerns are run using antibodies raised to purified w/t EPSPS from Brassica napus (expressed and purified using similar methods to those described above). In this case the strength of the cross reaction of the antibodies is less strong with rice EPSPS (or with endogenous plant wheat or maize EPSPS) than in the case that antibodies raised to rice EPSPS be used but still, nevertheless, provide sufficient reaction for useful quantitative information in relation to the standard curve to be derivable.

**EXAMPLE 20. Isolation of genomic DNA from transgenic plant material. PCR analysis. DNA probe preparation and hybridisation. Copy number and integrity of transgene.**

Genomic DNA is isolated from plants, plantlets and callus material using ,for example, the method of Dellporta et al (1983) in Chromosome Structure and Function: Impact of New Concepts, 18'th Stadler Genetics Symposium. J.P. Gustafson and R . Appels, eds). New York: Plenum Press) pp 263-282 or, alternatively, the DNAeasy kit (Qiagen) may be used. Transgenic callus and plant segregants that contain the mutated rice EPSPS transgene are identified using fluorescence PCR using oligonucleotide primers SEQ ID NO. 37 and 38 that are specific to the mutations within the rice EPSPS genomic sequence. The fluorescent dye SYBR green which intercalates with double stranded DNA, is included in the PCR so that samples containing the mutated rice EPSPS gene are detected by an increase in fluorescence in the sample which is detected using an ABI 3377 machine. Alternatively those skilled in the art will know that the primers may be fluorescently labelled and technologies such as molecular beacons and 'Scorpions' are available.

SEQ. ID. NO. 37

RiceDM Fwd2-3A 5' -gtg gaa cgc tgg aat tgc aat gca at -3'

SEQ. ID. NO. 38

Univeral Reverse 5' - gtt gca ttt cca cca gca gca gt - 3'

A typical PCR consist of , prepared in 96 well optical plates and sealed with Optical lids (PE Biosystems), is as follows in 25 µl total volume:

5.0 µl gDNA template (Qiagen Dneasy prep)

12.5 µl 2X SYBR Green premix  
2.5 µl 5pmol/µl stock forward primer  
2.5 µl 5pmol/ µl stock reverse primer  
2.5 µl dd H<sub>2</sub>O

5 The following cycling parameters are followed:

Stage 1: 50 C for 2 min  
Stage 2: 95 C for 10 min  
Stage 3: 95 C for 15 s  
60 C for 45 s

10 Changes in fluorescence within the samples are recorded every seven seconds from stage 3 of the reaction

For Southern blotting approximately 10 µg of DNA is used for each restriction digest. Genomic DNA is digested with suitable restriction enzymes (e.g Hind III) according to the manufacturer's instructions (e.g Promega). Restriction enzymes are chosen that cut both  
15 within and outside the mutant EPSPS sequence. DNA is separated using TAE (0.04M tris-acetate, 1 mM EDTA) 0.8% agarose gels. Southern blotting is carried out according to methods giving by Sambrook et al., 1989 using HyBond N+ nitrocellulose blotting membrane (AmershamPharmacia). The DNA is cross-linked to the membrane by exposure to UV illumination.

20 DNA fragments used for generating specific probes are isolated by purification on gels of restriction digests of plasmid DNA or generated by PCR. For example, a 700 bp fragment containing intron 1 of the rice EPSPS gene, is generated by PCR using primers as shown below.

SEQ. ID. NO 39

25 INT1/55' cccttcctcttgcgtaattccatttc 3'

SEQ. ID. NO. 40

INT1/35' gttgtgccccaataaccagag 3'

Such probes are labelled with <sup>32</sup>P using the random priming method, for example Ready-To-Go DNA labelling beads (AmershamPharmacia) and purified using, for example,  
30 MicroSpin G-25 columns (AmershamPharmacia).

Blots of DNA gels are prehybridized at 65 C in 5x SSC, 0.5% SDS, 2xDenhardt's solution, 0.15 mg/ ml denatured salmon sperm DNA for at least one hour. The blot is then

hybridized with denatured probe for 16-24 h at 65 C in fresh pre-hybridisation solution. Membranes are blotted dry and visualised using autoradiography.

Where Southern blotting indicates a single integration event of the transgene at a single locus, indicated by the probe hybridising with only a single specific sized restriction  
5 fragment, then the result is confirmed through a rehybridisation of the blot using an alternative probe. For controls, untransformed material is used. Additionally the blot may be probed further with hybridisation probes specific to other regions of the transgenic construct (for example the promoter, 5'UTR intron or upstream enhancer sequences) in order to verify the integrity of the construct. Additionally, in the case that *Agrobacterium* transformation is  
10 used, specific probes are used to indicate the presence or absence of any DNA extending from beyond the RB and LB of the super-binary vector.

## SEQ ID NO. 41 Rice EPSPS genomic (from ATG - WT sequence)

atggcggcgaccatggcgtccaacgccgcggctgcggcggcggtgtccctggaccaggccgtggcggcgctcggcggcggttctc  
gtcgcggaagcagctgcggctgcccgcgcggcgcgcggggggatgcgggtgcgggtgcgggcgckggggcggcgggaggcgg  
tgggtggcggcgtccgcgtcgtcgtcgtcggcggcagcgcggcgggcgaaggcggaggagatcgtgctccagcccatcaggagg  
5 atctccggggcggttcagctgccagggtccaagtcgctctccaacaggatcctcctcctcctccgcccctcctcagagggtgagacg  
cggatcccttctccttgcgtgaattccatttcttgagatgagattttagggggtttattaggtgaggtggctgtgtttgtgaa  
atcctaggaattatctctcaagtcaatctaacgatgagatataactgaggttctgggttttaatcacacactcatataaccaat  
ttattgaaacattttgggttggcataagaaactgcttacgaaggtatgatatcctcctacatgtcaggtactaaattttcac  
gacgggtatgatccactcaaaacaagtttcttaacgagctcgttgagggtctgttatgaaatttgtgtaaaactaaggcaactttg  
10 gaggtttcgcactgtaccaatgttatgtttgaacattttgcaagcagtgctttctccaaaattatgcaattttgagggtcct  
ctacatcattataattccccaatacattgctctttattcttaatagctttgatcgcgaaatttaacatttttaattcttgagct  
gttatttttagcatcagtttatcatgagccatgtttgggtactaaatatacaatcccttgggtttatttgtttccaagcatgt  
cattaacttatcttaatgtggacaagaaactgatgcctgcttacattgctattatttcaagcgggtattgatcctttgacatg  
tgattgatcatttttttttctctggttattagggcacaacagtggtggacaacttgctgaacagtgaggatgttactacatg  
15 cttgaggccctgaaagccctcgggctctctgtggaagcagataaagttgcaaaaagagctgtagtcgttggctgtgggtggcaa  
gtttcctgttgagaaggatgcgaaagaggaagtgaactcttcttggggaacgctggaactgcaatgcgaccattgacagcag  
ccgtgactgctgctgggtggaaatgcaacgtatgttttttttttaattgtttatgaaaatatgtatggaattcatgggggtatgt  
tttatgaccttttctttaccatcagttatgtgcttgatggagtgccacgaatgaggagagaccgattgggtgacttgggtgt  
cgggttgaaacaacttgggtgcggatgtcgactgttcttggcactgaatgccacctgttcgtgtcaaggggaattggaggac  
20 ttctcgtggcgaagggttagttactcctaaactgcaccttctgtacttctgtatgcacctcaattcttctgcaaccttctgcat  
ttataaggaacattctatgatgcaattcgaccttacactgcacagtaacttgaaatgtttcatgcttaataatgccatat  
tcttgccaagctcaagcgagcaatatttgtttgaatttgggtaccatattttctgtatatttgggcattccttttgggtcttgat  
gtcttcttttgaattagcatttaactgaattacactcaacaggttaagctctctggttccatcagcagtcagttacttgagtgc  
cttgctgatggctgctccttggcccttggggatgtggagatcgaaatcattgacaaaactaatctccattccttacgttgaaa  
25 tgacattgagattgatggagcgttttgggtgtaaggcagagcattctgatagttgggacagattctatatgaggggaggcag  
aagtacaagtaagcttctacctgccttactgagctgaattattcgggtgtttatgattaactccctaaactaacccttttct  
tttcttggcattgacagatctcctggaaatgcctatgttgaaagtgatgcctcaagcgcgagctatttcttggctgggtgctg  
caatcactggaggcactgtgacagttcaaggttgtggtagcaccagtttgacaggtataactgtagtgctgttttgacattct  
accgttttagtcaagtttagtcagtagtcacatatcagaatatagcacaatctgtattatgccactgttaatacaaatagcctt  
30 gacctagagagtgctatataccctagcttaactctcaaaactaaacagttctcttgtggcttgctgtgctgttatgttccctga  
cctacatgttaataattacagggtgatgtcaaatttgctgaggtacttgagatgatgggagcaagggttacatggactgacacc  
agtgtaacctgaactgggtccaccacgtgagccttatgggaagaaacacctgaaagctgttgatgtcaacatgaacaaaatgcc  
tgatgttgccatgaccttggcgttgttgactcttcgctgatgggtccaactgctatcagagatgggtaaacattaaaggcctat  
tatacctgttctatcatactagcaattactgcttagcattgtgacaaaacaaataaccaaactttcttcaaaaataacttagaa  
35 atataagaaagggttcgttttgtgtggtaaacagtactactgtagtttcagctatgaagtttgctgctggcaattttctgaacg  
gtttcagctaaattgcatgtttgttcatcatacttatccattgtctccacagtggttccctggagagtaaaaggaaaccgaaa  
ggatgggtgcaattcggaccgagctaacaaaggtaaatcattaggtcccgtgtcctttcattcttcaagtagtttgttcata  
agttgaattctccttcaatgatgtttaaattcatcatcttcttttttgggtgttggtgccagctgggagcatcgggtgaagaagg  
tcttgactactgcatcatcaccacacgggagaagctgaacatcacggcaatcgacacctacgatgatcacaggatggccatgg  
40 ccttctcctcgtgctgctgcgcgacgtgcccgtgacgatcagggaacctgggtgcacccgcaagaccttcccaactacttc  
gacgttctaagcactttcgtcaggaactgaactgagcttttaaaagagtgaggtctaggttctgttgtctgtctgtccatcat  
ccatgtgttgactgttgagggaactcgtttcttcttcttcttcacgagatgagtttttggtgtgctgttaatactagtttgtgac  
aaaggctgcgttacataagggtgatgagaattgaggtaaaatgagatctgtacactaaattcattcagactgttttggcataaa  
gaataatttggccttctgcgatttcagaagctataaattgccatctcactaaattctccttgggtcctcatggcaatgcaacga  
45 cagtggtgaagcactgaagcccgtaatgctctatcaccacatgtacgacagaaccatatatgtccatattgtacaactcgagtg  
ttgtttgagtgccagcaaaactgggtgaccaagccacacgagagagaataactataaactcaatcatacataacaagcccaagc  
aacattagacagaacacaacaacactcg

## Rice EPSPS genomic (from ATG including double mutant - shown)

ggcgtccaacgccgcggctgcggcggcggtgtccctggaccaggccgtggcggcgctcggcggcggttctc  
tgcggctgccgcgcgcggcgcgcgggggatgcgggtgcgggtgcgggcgckggggcgggaggcg  
gcgtcgtcgtcgtcgtcggcggcagcgcggcggaaggcggaggagatcgtgctccagcccatcaggag  
tcagctgccagggtccaagtcgctctccaacaggatcctcctcctcctccgccctcctccgagggtgagacg  
ttgcgtgaattccatttctggagatgagattttagggggtttattagggtgagggtggctgtgtttgtgaa  
ctctcaagtcgaatctaacgatgagatataactgaggttctgggttttaatacacacactcatataaccaat  
tgggttggcataagaaactgcttacgaaggatgatctcctctacatgacaggctactaaattttcac  
ctcaaaacaagtttcttaacgagtcgtggtgaggtctgttatgaaatttgtgtaactaaggcaactttg  
taccaatgttatgtttgaacattttgcaagcagtgctttctcccaaaattatgcaattttgaggctcct  
ttccccaatacattgctctttattcttaataagctttgatcgcgaaatttaacatttttaattcttgagct  
tcagtttatcatgagccatgtttgggtactaaatatacaatcccttgggtttatttgtttccaagcatgt  
aatgtggacaagaaactgatgcctgcttacattgctattatttcaagcgggtattgatcctttgacatg  
cttttctctggttattagggcacaacagtggtggacaacttgctgaacagtgaggatgttcaactacatg  
agccctcgggctctctgtggaagcagataaagtgcacaaaagagctgtagtcgttggctgtggtggcaa  
aggatgcgaaagaggaagtgcacactctcttggggaacgctggaaTtgcaatgcgaTcatgacagcag  
ggtggaaatgcaacgtatgttttttttttaagtgttatgaaaatgtatggaattcatggggtatgt  
ctttaccatcagttatgtgcttgatggagtgccacgaatgaggagagagaccgattggtgacttgggtgt  
ctgggtgcggatgtcgactgtttccttggcactgaatgccacctgttcggtgtcaagggaattggaggac  
gttagttactcctaaactgcacctcttctgacttctgtatgcacctaattctttgtcaaccttctgcat  
ctatgatgcaattcgacctacactgcacagtaactgaaatgtttcatgcttaataatgccaat  
tgcgagcaatatttgtttgaatttgggtaccatattttgtatatttgggcattccttttgggtcttgat  
agcatttaactgaattacactcaacagggttaagctctctggttccatcagcagtcagtaacttgagtgc  
ctccttggcccttggggatgtggagatcgaaatcattgacaaactaatctccattccttacgttgaaa  
tggagcgttttgggtgtgaaggcagagcatctgatagttgggacagattctatattaaggaggggcag  
tctacctgccttactgagctgaattattcgggtgtttatgattaactccttaactaaccctttttct  
cagatctcctggaaatgcctatgttgaagggtgatgcctcaagcgcgagctatttcttggctggtgctg  
ctgtgacagttcaagggtgtggtacgaccagtttgacaggtataactgtagtgctgttttgacattct  
ttagtcagtagtcacataattcagaatatagcacaatctgtattatgccactgttaatacaatacgtt  
atataccctagcttaattctcaactaaacagttctcttgtggcttgcctgtgctgttatgttccctga  
tcacagggtgatgtcaaatttgctgaggtacttgagatgatgggagcaaagggtacatggactgacacc  
gggtccaccacgtgagccttatgggaagaaacacctgaaagctgttgatgtcaacatgaacaaaatgcc  
ccttgcggttgttgactcttcgctgatgggtccaactgctatcagagatggtaaacatttaaggcctat  
atactagcaattactgcttagcatgttgacaaaacaaataaccaaactttcttcaaaaataacttagaa  
gttttgtgtggtaaacagtaactactgtagtttcagctatgaagtttgcctgtggcaattttctgaacg  
catgtttgttcatcatacttatccattgtcttcacagtggttccctggagagtaaaaggaaaccgaaa  
ggaccgagctaaacaaaggtaaatcattaggtcccggtgtcctttcattcttcaagtagtttgttcata  
caatgatgtttaaattcatcatcttcttttttgggtgtgtgcccagctgggagcatcgggtgaagaagg  
catcaccaccgggagaagctgaacatcacggcaatcgacacctacgatgatcacaggatggccatgg  
ctgcgcgcgacgtgccggtgacgatcagggaccctgggtgcaccgcgaagaccttcccaactacttc  
ttcgtcaggaactgaactgagcttttaaaagagtgaggtctaggttctgtgtgtctgtctgtccatcat  
tgagggaactcgtttcttcttttcttcacgagatgagtttttgtgtgcctgtaatactagtttgtagc  
taagggtgatgagaattgaggtaaaatgagatctgtacactaaattcattcagactgttttggcataaa  
tgcgatttcagaagctataaattgccatctcactaaattctccttgggtcctcatggcaatgcaacga  
tagcccgtaatgctctatcaccacatgtacgacagaacatataatgtccatgtgacaactcgagtg  
tcaaaactggctgaccaagccacacgagagagaatactataaactcaatcatataacaagcccaagc  
caacaacactcg

## SEQ ID NO. 43 Maize PolyU Enhancer

ttcagccttcgatgtggatgcaacagcttcacaggattccattaaatcgtagccattgtgtcaaagtttgctttgccaacggt  
atctattttatttttagaaaaccagctttgaccagccgcctctttacgtttggcacaatttagctgaatccggcgccatgg  
5 caaggtagactgcagtgcagcgtgacccggtcgtgccctctctagagataatgagcattgcatgtctaagttataaaaaatt  
accacatatattttttgtcacacttggttgaagtgcagtttatctatctttatacatatatttaaactttactctacgaataat  
ataatctatagtactacaataatatcagtgttttagagaatcatataaatgaacagttagacatggcttaaaggacaattgag  
tattttgacaacaggactctacagttttatcttttagtgtgcatgtgttctcttttttttgcataatagcttcacctata  
taatacttcatccattttattagtagacatccatttaggggttaggggttaatgggttttatagactaatttttttagtagacatcta  
10 ttttattctatttttagcctctaaattaagaaaaactaaaactctatttttagtttttttatttaataatttagatataaaataga  
ataaaataaagtactaaaaatataacaaaataccctttaagaaataaaaaaactaaggaaacatttttctgtttcgagtag  
ataatgccagcctgttaaacgccgtcgacgagtcacggacaccaaccagcgaaccagcagcgtcgcgctcgggccaagcgaa  
gcgagcggcgccgcatctctgtcgtgcctctggaccct

## 15 SEQ ID NO. 44 Rice Actin Enhancer

gatatccctcagccgctttcactatctttttgcccagagtcattgtcatgtgaaccttggcatgtataatcgggtgaattgcg  
tcgattttctcttataggtgggccaatgaatccgtgtgatcgcgctgattggctagagatatgtttctctcttgttggatg  
tattttcacataatcatatgcatacaaatatttcattacactttatagaaatgggtcagtaataaaccttatcactatgtct  
gggtgtttcattttatttgcttttaaacgaaaattgacttctgattcaatatttaaggatcgtaacgggtgtgcagttactaa  
20 attctgggttgtaggaactatagtaaaactattcaagtcttcaactatttgtgcactcacctctcgccacatcaccacagatgtt  
attcacgtcttaaatgtgaactacacatcatattgacacaatatttttttaataagcgattaaaacctagcctctatgtca  
acaatgggtgtacataaccagcgaagtttagggagtaaaaaacatcgccctacacaaagttcgctttaaaaaataaagagttaa  
ttttactttggaccacccttcaaccaatgtttcactttagaacgagtaattttattattgtcactttggaccaccctcaaatc  
ttttttccatctacatccaatttatcatgtcaaagaaatgggtctacatacagctaaggagatttatcgacgaatagtagctag  
25 catactcgaggtcatctatgtcttgagaagagagtcgggatagtcacaaaataaaacaaaggtaagattacctgggtcaaaagt  
gaaaacatcagttaaaagggtggtataaagtataaatatcggtataaaagggtggcccaaagtgaaatttactctttctactat  
tataaaaaattgaggtatgtttttgtcggtagctttgatacgtcatttttgatgaattgggttttaagtatttctgcttttgga  
atgcatactctgtattttgagtcgggttttaagtctggtttgcttttgtaatacagagggatttgatataagaaatatcttataaa  
aaaccatagctaatgtgacataatttttgagaaaaatatattcaggcgaattctcacaaatgaacaataaagattaaa  
30 atagctttcccccggtgcagcgcaggggtattttttctagtataaaataaaagataaaacttagactcaaaacatttacaaaaac  
aaccctaaagttcctaaagcccaaagtgctatccacgatccatagcaagcccagcccaaccaaccaaccaacccacccc  
agtccagccaactggacaatagtcctccacacccccccactatcacctgaggtgtccgcacgcacgcacgtctcgagccaa  
aaaaaaaaaagaaagaaaaaaaagaaaaagaaaaaacagcaggtgggtccgggtcggtggggccggaacgcgaggaggatc  
gcgagca

35

## SEQ ID NO. 45 Rice Genomic G1 EPSPS (to ATG)

gttgggtgggtgagagtgcagacaccgacggaacggaaggagaaccacgcccgttggatttttcttttttaccttttcaaat  
aatttaaaaaataaaaccatttttaaaaacttatcttcaaatacaaatcttttaaaaacactaacacgtgacacacagcgggca  
cgtcacccaaacggcggtgacaatatgtttttgccacaccaatccagctgggtgtggacaaaatgttcatatattgaaaataaa  
40 atttaaaacaatttatattttttatctatatcattataaaaaattgaagatgtttttaccgggtattttgttactcatttgtgca  
tgagtcgggttttaagtgtgttcgcttttggaatacatatccgtatttgagtatgttttaagttcggttcggtttttgaaat  
acaaaaggaatcgtaaaataaatctatttttaaaaactcgcatgctaacttgagacgatcgaactgctaattgcagctcataa  
ttttccaaaaaaaatatatccaaacgagttcttatagtagatttcaccttaattaaaacataataatgttaccgggtacaa  
cgcacgagttttttataagtaaaattaaaagtttaaaataaataaaaatcccgccaccacggcgcgatggtaaaagggggac  
45 gcttctaaacggggccgggcacgggacgatcgggcccggaaccggcccatctaaccgctgtaggccaccggccaccaatccaa



ctccgtactacgtgaagcgcgtggatccgcaaccgcgttaagcagtcacacgactcgactcgactcgcgactcgccgtggtag  
gtggcaacccttcttctctctctatttcttcttcttctctctctctcgcctcaccacaccaaccgcaccaaccccaaccccg  
cgcgcgctctccccctctccccctcccaccaacccccaccccatcctcccgacctccacgcgcgcggaatg

5 SEQ ID NO. 46 Rice Genomic G3 EPSPS (to ATG)

ttaattaaaacatataaatgttcacccggtacaacgcacgagtatTTTTATAAGTAAAATTAAAAGTTAAAATAAATAAAAA  
tcccgccaccacggcgcgatggtaaaagggggacgcttctaaccgggcccgggcacgggacgatcgccccgaaccggcccat  
ctaaccgctgtagggccaccgcccaccaatccaactcgcactacgtgaagcgcgtggatccgcaaccgcgttaagcagtcacac  
cgactcgactcgactcgcgactcgccgtggtaggtggcaacccttcttctctctctatttcttcttcttctctctctctccg  
10 cctcaccacaccaaccgcaccaacccccaccccgcgcgcgctctccccctctccccctcccaccaacccccaccccatcctccga  
cctccacgcgcgcggaatg

SEQ ID NO. 47 Rice Genomic G2 EPSPS +Maize Adh1 intron

gccacaccaatccagctggtgtggacaaaatgttcatatattgaaaataaaaatttaaaacaatttatattttttatctatct  
15 attataaaaattgaagatgtttttaccggtattttgttactcatttgtgcatgagtcgggtttttaagttgttcgcttttgga  
aatacatatccgtatttgagtagtgtttttaagttcggttcttttgaaatacaaaaaggaatcgtaaaaataatctattttaa  
aaaactcgcatgctaacttgagacgatcgaaactgctaattgcagctcataattttccaaaaaaaatatatccaaacgagttc  
ttatagtagattttcaccttaattaaaacatataaatgttcacccggtacaacgcacgagtatTTTTATAAGTAAAATTAAAAG  
tttaaaataaaataaaaaatcccgccaccacggcgcgatggtaaaagggggacgcttctaaccgggcccgggcacgggacgatcgg  
20 ccccgaaaccggcccatctaaccgctgtagggccaccgcccaccaatccaactcgcactacgtgaagcgcgtggatccgcaac  
cgcgttaagcagtcacacgactcgactcgactcgcgactcgccgtggtaggtggcaacccttcttctctctctatttcttct  
tcttctctctcttctccgctcaccacaccaaccgcaccaacccccaccccgcgcgcgctctccccctctccccctcccaccaacc  
ccaccccatcctcccgacctccacgcgcgcgcgagatcaagtgcagggtccgcttgtttctctctctgtctcttgatctga  
ctaactcttggtttatgattcgttgagtaattttggggaaagctagcttcggtccacagttttttttcgatgaacagtgccgca  
25 gtggcgctgatcttgatgctatcctgcaatcgtgggtaacttatttcttttatatccttcaactcccatgaaaaggctagtaa  
tctttctcgatgtaacatcggtccagcactgctattaccgttggtccatccgacagtcgtggtgaacacatcatacgatatg  
agcaaagatctatcctccctgttctttaatgaaagacgtcattttcatcagtatgatctaagaatgttgcaacttgcaaggag  
gcgtttctttctttgaatttaactaactcgttgagtggccctgtttctcggaagtaaggcctttgctgctccacacatgtcca  
ttcgaattttaccgtgttagcaagagcgaaaagttgcatcttgatgatttagcttgactatgcgattgctttctctggaccc  
30 gtgcagctgcggatg

SEQ ID No. 48 Maize Adh1 intron

gtccgccttggtttctctctgtctcttgatctgactaatcttggtttatgattcgttgagtaattttggggaaagctagcttc  
gtccacagtttttttttcgatgaacagtgccgcagtgccgctgatcttgatgctatcctgcaatcgtgggtaacttatctct  
35 tttatataccttcaactcccatgaaaaggctagtaatttttctcgatgtaacatcggtccagcactgctattaccgtgtggtccat  
ccgacagtcgtggtgaacacatcatacgatattgagcaaagatctatcctccctgttctttaatgaaagacgtcattttcatc  
agtatgatctaagaatgttgcaacttgcaaggaggcgtttcttctttgaatttaactaactcgttgagtggccctgtttctc  
ggacgtaaggcctttgctgctccacacatgtccattcgaattttaccgtgttagcaagagcgaaaagttgcatcttgatga  
tttagcttgactatgcgattgctttctctggacccgtgcag

40

**CLAIMS**

1. An isolated polynucleotide comprising the sequence depicted in SEQ ID No. 41.
- 5 2. A polynucleotide encoding an EPSPS, excluding the cDNA encoding the rice and corn EPSPS, which polynucleotide is complementary to one which when incubated at a temperature of between 65 and 70°C in 0.1 strength citrate buffered saline containing 0.1% SDS followed by rinsing at the same temperature with 0.1 strength citrate buffered saline containing 0.1% SDS still hybridises with the sequence  
10 depicted in SEQ ID No. 41.
3. A polynucleotide encoding an EPSPS obtainable by screening plant genomic DNA libraries with a polynucleotide constituting an intron within the SEQ ID No. 41  
15 sequence.
4. An isolated polynucleotide comprising a region encoding a chloroplast transit peptide and a glyphosate resistant 5-enolpyruvylshikimate phosphate synthase (EPSPS) 3' of the peptide, the said region being under expression control of a plant operable promoter, with the *provisos* that the said promoter is not heterologous with respect to  
20 the said region, and the chloroplast transit peptide is not heterologous with respect to the said synthase.
5. A polynucleotide according to any one of claims 1 to 4, comprising the following components in the 5' to 3' direction of transcription:-  
25 (i) At least one transcriptional enhancer being that enhancing region which is upstream from the transcriptional start of the sequence from which the enhancer is obtained and which enhancer *per se* does not function as a promoter either in the sequence in which it is endogenously comprised or when present heterologously as part of a construct;  
30 (ii) The promoter from the rice EPSPS gene;  
(iii) The rice genomic sequence which encodes the rice EPSPS chloroplast transit peptide;

(iv) The genomic sequence which encodes the rice EPSPS;

(v) A transcriptional terminator;

wherein the rice EPSPS coding sequence is modified in that a first position is mutated so that the residue at this position is Ile rather than Thr and a second position is mutated so that the residue at this position is Ser rather than Pro, the mutations being introduced into EPSPS sequences which comprise the following conserved region GNAGTAMRPLTAAV in the wild type enzyme such that modified sequence reads GNAGIAMRSLTAAV.

- 5
6. A polynucleotide according to claim 5, wherein the said enhancer comprises a sequence, the 3' end of which is at least 40 nucleotides upstream of the closest transcriptional start of the sequence from which the enhancer is obtained.
- 10
7. A polynucleotide according to either of claim 5 or 6, wherein the enhancer comprises a region the 3' end of which is at least 60 nucleotides upstream of the said closest start.
- 15
8. A polynucleotide according to claim 5, wherein the said enhancer comprises a sequence the 3' end of which is at least 10 nucleotides upstream from the first nucleotide of the TATA consensus of the sequence from which the enhancer is obtained.
- 20
9. A polynucleotide according to any one of claims 1-8, comprising first and second transcriptional enhancers.
- 25
10. A polynucleotide according to claim 9, wherein the first and second enhancers are tandemly present in the polynucleotide.
- 30
11. A polynucleotide according to any one of claims 1-10, wherein the 3' end of the enhancer, or first enhancer, is between about 100 to about 1000 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.

12. A polynucleotide according to any one of claims 1-11, wherein the 3' end of the enhancer, or first enhancer, is between about 150 to about 1000 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.
13. A polynucleotide according to any one of claims 1-12, wherein the 3' end of the enhancer, or first enhancer, is between about 300 to about 950 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.
14. A polynucleotide according to any one of claims 1-13, wherein the 3' end of the enhancer, or first enhancer, is between about 770 and about 790 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.
15. A polynucleotide according to any one of claims 1-13, wherein the 3' end of the enhancer, or first enhancer, is between about 300 and about 380 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.
16. A polynucleotide according to any one of claims 1-13 and 15, wherein the 3' end of the enhancer, or first enhancer, is between about 320 and about 350 nucleotides upstream of the codon corresponding to the translational start of the EPSPS transit peptide, or the first nucleotide of an intron in the 5' untranslated region.
17. A polynucleotide according to any one of claims 1-16, wherein the region upstream of the promoter from the rice EPSPS gene comprises at least one enhancer derived from a sequence which is upstream from the transcriptional start of either the maize polyubiquitin or rice actin promoters.

18. A polynucleotide according to claim 17, comprising in the 5' to 3' direction a first enhancer comprising a transcriptional enhancing region derived from a sequence which is upstream from the transcriptional start of the rice actin promoter and a second enhancer comprising a transcriptional enhancing region derived from a sequence which is upstream from the transcriptional start of a rice actin promoter.
19. A polynucleotide according to any one of claims 1-18, wherein the nucleotides 5' of the codon which constitutes the translational start of the rice EPSPS chloroplast transit peptide are Kozak preferred.
20. A polynucleotide according to any one of claims 1-19, wherein 5' of the rice genomic sequence which encodes the rice EPSPS chloroplast transit peptide there is located a non-translated region which comprises a sequence which functions as an intron.
21. A polynucleotide according to claim 20, wherein the non-translated region comprises an intron, wherein the intron is the maize ADHI intron.
22. A polynucleotide according to either of claims 21 or 20, wherein the non-translated region comprises the sequence depicted in SEQ ID NO. 48.
23. A polynucleotide according to claim 22, wherein the non-translated region comprises an intron, wherein the intron is a rice actin intron 1 or a maize polyubiquitin intron.
24. A polynucleotide according to any one of claims 1-23, which comprises a virally derived translational enhancer located within the non translated region 5' of the rice genomic sequence which encodes the rice EPSPS chloroplast transit peptide.
25. A polynucleotide according to any one of claims 1-24, further comprising regions encoding proteins capable of conferring upon plant material containing it at least one of the following agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, dessication, and herbicides.

26. A polynucleotide according to claim 25, wherein the herbicide is other than glyphosate.
- 5 27. A polynucleotide according to either of claims 25 or 26, wherein the insect resistance conferring regions encode crystal toxins derived from Bt, including secreted Bt toxins; protease inhibitors, lectins, Xenhorabdus/Photorhabdus toxins; the fungus resistance conferring regions are selected from the group consisting of those encoding known AFPs, defensins, chitinases, glucanases, Avr-Cf9; the bacterial resistance
- 10 conferring regions are selected from the group consisting of those encoding cecropins and techyplesin and analogues thereof; the virus resistance regions are selected from the group consisting of genes encoding virus coat proteins, movement proteins, viral replicases, and antisense and ribozyme sequences which are known to provide for virus resistance; the stress, salt, and drought resistance conferring regions are selected
- 15 from those that encode Glutathione-S-transferase and peroxidase, the sequence which constitutes the known CBF1 regulatory sequence and genes which are known to provide for accumulation of trehalose.
28. A polynucleotide according to claim 27, wherein the insect resistance conferring
- 20 regions are selected from the group consisting of cryIAC, cryIAB, cry3A, Vip 1A, Vip 1B, cystein protease inhibitor, and snowdrop lectin genes.
29. A polynucleotide according to any preceding claim, which is modified in that mRNA
- 25 instability motifs and/or unwanted splice regions are removed, or crop preferred codons are used so that expression of the thus modified polynucleotide in a plant yields substantially similar protein having a substantially similar activity/function to that obtained by expression of the protein encoding regions of the unmodified polynucleotide in the organism in which they are endogenous.
- 30 30. A polynucleotide according to the preceding claim, wherein the degree of identity between the modified polynucleotide and a polynucleotide endogenously contained

within the said plant and encoding substantially the same protein is such as to prevent co-suppression between the modified and endogenous sequences.

- 5           31.    A polynucleotide according to the preceding claim, wherein the said degree is less than about 70%.
32.    A vector comprising the polynucleotide of any preceding claim.
- 10          33.    Plant material which has been transformed with the polynucleotide of any one of claims 1-31, or the vector of claim 32.
34.    Plant material which has been transformed with the polynucleotide of any one of claims 1-31 or the vector of claim 32, and which has been, or is, further transformed with a polynucleotide comprising regions encoding proteins capable of conferring  
15           upon plant material containing it at least one of the following agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, desiccation, and herbicides.
35.    Morphologically normal, fertile whole plants which have been regenerated from the  
20           material according to either of claims 33 or 34, their progeny seeds and parts.
36.    Morphologically normal fertile whole plants which comprise the polynucleotide of any one of claims 1-31 and which result from the crossing of plants which have been regenerated from material transformed with the polynucleotide of any one of claims  
25           1-31 or the vector of claim 32, and plants which have been transformed with a polynucleotide comprising regions encoding proteins capable of conferring upon plant material containing it at least one of the following agronomically desirable traits: resistance to insects, fungi, viruses, bacteria, nematodes, stress, desiccation, and herbicides, the progeny of the resultant plants, their seeds and parts.  
30
37.    Plants according to either of claims 35 or 36, selected from the group consisting of field crops, fruits and vegetables such as canola, sunflower, tobacco, sugar beet,

5 cotton, maize, wheat, barley, rice, sorghum, tomato, mango, peach, apple, pear, strawberry, banana, melon, potato, carrot, lettuce, cabbage, onion, soya spp, sugar cane, pea, field beans, poplar, grape, citrus, alfalfa, rye, oats, turf and forage grasses, flax and oilseed rape, and nut producing plants insofar as they are not already specifically mentioned, their progeny, seeds and parts.

38. Maize plants according to any one of claims 35-37.
- 10 39. A method of selectively controlling weeds in a field, the field comprising weeds and plants or progeny according to any one of claims 35-38, the method comprising application to the field of a glyphosate type herbicide in an amount sufficient to control the weeds without substantially affecting the plants.
- 15 40. A method according to the preceding claim, further comprising application to the field either before or after application of the glyphosate herbicide of one or more of the following: a herbicide, insecticide, fungicide, nematocide, bacteriocide and an anti-viral.
- 20 41. A method of producing plants which are substantially tolerant or substantially resistant to glyphosate herbicide, comprising the steps of:  
(i) transforming plant material with the polynucleotide of any one of claims 1 to 31 or the vector of claim 32;  
(ii) selecting the thus transformed material; and  
(iii) regenerating the thus selected material into morphologically normal fertile whole plants.
- 25 42. A method according to the preceding claim, wherein the transformation involves the introduction of the polynucleotide into the material by: (i) biolistic bombardment of the material with particles coated with the polynucleotide; or (ii) impalement of the material on silicon carbide fibres which are coated with a solution comprising the polynucleotide; or (iii) introduction of the polynucleotide or vector into
- 30



Agrobacterium and co-cultivation of the thus transformed Agrobacterium with plant material which is thereby transformed and is subsequently regenerated.

43. A method according to the preceding claim, wherein the transformed material is  
5 selected by its resistance to glyphosate.
44. Use of the polynucleotide of any one of claims 1 to 31, or the vector of claim 32, in  
the production of plant tissues and/or morphologically normal fertile whole plants  
which are substantially tolerant or substantially resistant to glyphosate herbicide.  
10
45. Use of the polynucleotide of any one of claims 1 to 31, or the vector of claim 32, in  
the production of a herbicidal target for the high throughput in vitro screening of  
potential herbicides.
- 15 46. A method of selecting biological material transformed so as to express a gene of  
interest, wherein the transformed material comprises the polynucleotide of any one of  
claims 1 to 31 or the vector of claim 32 and wherein the selection comprises exposing  
the transformed material to glyphosate or a salt thereof, and selecting surviving  
material.
- 20 47. A method according to the preceding claim, wherein the biological material is of  
plant origin.
48. A method according to the preceding claim, wherein the plant is a monocot.
- 25 49. A method according to the preceding claim, wherein the monocot is selected from  
group consisting of barley, wheat, corn, rice, oats, rye, sorghum, pineapple, sugar  
cane, banana, onion, asparagus, leek,
- 30 50. A method for regenerating a fertile transformed plant to contain foreign DNA  
comprising the steps of:  
(a) producing regenerable tissue from said plant to be transformed;

- (b) transforming said regenerable tissue with said foreign DNA, wherein said foreign DNA comprises a selectable DNA sequence, wherein said sequence functions in a regenerable tissue as a selection device;
- (c) between about one day to about 60 days after step (b), placing said  
5 regenerable tissue from step (b) in a medium capable of producing shoots from said tissue, wherein said medium further contains a compound used to select regenerable tissue containing said selectable DNA sequence to allow identification or selection of the transformed regenerated tissue;
- (d) after at least one shoot has formed from the selected tissue of step (c)  
10 transferring said shoot to a second medium capable of producing roots from said shoot to produce a plantlet, wherein the second medium optionally contains the said compound; and
- (e) growing said plantlet into a fertile transgenic plant wherein the foreign DNA is transmitted to progeny plants in Mendelian fashion, wherein between step (b) and  
15 step (c) there is an optional step of placing the transformed material onto callus inducing medium, characterised in that the foreign DNA is, or the selectable DNA sequence comprised by the foreign DNA comprises, the polynucleotide according to any one of claims 1 to 31 or vector of claim 32, and the said compound is glyphosate or a salt thereof.
- 20
51. A method according to the preceding claim, wherein the plant is a monocot selected from the group consisting of banana, wheat, rice, corn and barley.
52. A method according to either of claims 50 or 51, wherein the said regenerable tissue  
25 is selected from the group consisting of embryogenic calli, somatic embryos, immature embryos etc.




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 7




Figure 6






Figure 8




Figure 9




Figure 11




Figure 10




Figure 12




Figure 13




Figure 15

Figure 14




Figure 16




Figure 17a






Figure 17b




Figure 17c

I

## SEQUENCE LISTING

&lt;110&gt; Zeneca Ltd

&lt;120&gt; Herbicide Resistant Plants

&lt;130&gt; ppd50450

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 48

&lt;170&gt; PatentIn Ver. 2.0

&lt;210&gt; 1

&lt;211&gt; 32

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 1

gcacargcig caagigaraa igccatigcc at

32

&lt;210&gt; 2

&lt;211&gt; 29

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 2

gcwggaacwg cmatgcgicc rytiacigc

29

&lt;210&gt; 3

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 3

attttttttt cttctctcct tctccgctc

30

&lt;210&gt; 4

&lt;211&gt; 38

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 4

gagctccccg ggcgagtgtt gttgtgttct gtctaata

38

&lt;210&gt; 5

&lt;211&gt; 36

<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 5  
gcttacgaag gtatgatatc ctctacatg tcaggc 36

<210> 6  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 6  
gcagtcacgg ctgctgtcaa tgatcgcat t gcaattccag cgttcc 46

<210> 7  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 7  
ggaacgctgg aattgcaatg cgatcattga cagcagccgt gactgc 46

<210> 8  
<211> 55  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 8  
ggtgggcatt cagtgccaa gaaacagtcg acatccgcac caagttgttt caacc 55

<210> 9  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 9  
cgctgcagc tcgaggttgg ttggtgagag tgagacacc 39

<210> 10  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 10  
 cgcctgcagc tcgaggccac accaatccag ctggtgtgg 39  
  
 <210> 11  
 <211> 22  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:primer  
  
 <400> 11  
 gaacctcagt tatactctcat cg 22  
  
 <210> 12  
 <211> 43  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:primer  
  
 <400> 12  
 gcggccgcac tagtggccgg ccatcagcgg ccagcttttg ttc 43  
  
 <210> 13  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:primer  
  
 <400> 13  
 ttaactagtg aggaggccgc ctgccgtgc 29  
  
 <210> 14  
 <211> 45  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:primer  
  
 <400> 14  
 cgcctctaga ggccggccga tatccctcag ccgcctttca ctatc 45  
  
 <210> 15  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220>  
 <223> Description of Artificial Sequence:primer  
  
 <400> 15  
 cgctgcagtg ctgcgatcc tctcgtttt tcc 33  
  
 <210> 16  
 <211> 36  
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 16

gcgctgcagg atatccctca gccgcctttc actatc

36

<210> 17

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 17

gcgttaatta atgctcgcga tcctcctcgc ttttcc

36

<210> 18

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 18

cccacctcc cgacctccac gccgccggca ggatcaagtg caaaggccg ccttgtttct 60  
cctctg 66

<210> 19

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 19

gacgccatgg tcgccgccat ccgcagctgc acgggtccag gaaagcaatc

50

<210> 20

<211> 35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 20

cgagttctta tagtagattt caccttaatt aaaac

35

<210> 21

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 21  
ggacccgtgc agctgcggta ccatggcggc gaccatggc 39

<210> 22  
<211> 39  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 22  
gccatggctc cgcctatggt accgcagctg cacgggtcc 39

<210> 23  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 23  
tctctagact cagccgcctt tcactac 27

<210> 24  
<211> 42  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 24  
aaacccgggt ttggaagcgg agggaggaag gaggagataa ag 42

<210> 25  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 25  
accctcccct ctctaaatcg attggtggga ggggagag 38

<210> 26  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 26  
ggtctaccta caaaaaagct ccgcacgagg gtaccgccgc tggtac 46

<210> 27  
<211> 34  
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 27  
ccttcgcctc ccctccttcc tectctatatt cttc 34

<210> 28  
<211> 33  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 28  
gttggtggga ggggagagat ttagctaacc acc 33

<210> 29  
<211> 42  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 29  
gttttttcga ggcgtgctcc catggcggcg accatggcgt cc 42

<210> 30  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 30  
ggaggatatc ataccttcgt aagc 24

<210> 31  
<211> 40  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 31  
gcgctcgagt cagttcctga cgaaagtgct tagaacgtcg 40

<210> 32  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 32



gcgcatatga aggcggagga gatcgtgc 28

<210> 33  
<211> 28  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 33  
gcgcatatga aggcggagga gatcgtgc 28

<210> 34  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 34  
gcagtcacgg ctgctgtcaa tgatcgatt gcaattccag cgttcc 46

<210> 35  
<211> 40  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 35  
gcgctcgagt cagttcctga cgaaagtgt tagaacgtcg 40

<210> 36  
<211> 46  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 36  
ggaacgctgg aattgcaatg cgatcattga cagcagccgt gactgc 46

<210> 37  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:primer

<400> 37  
gtggaacgct ggaattgcaa tgcaat 26

<210> 38  
<211> 23  
<212> DNA  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 38

gttgcatttc caccagcagc agt

23

&lt;210&gt; 39

&lt;211&gt; 27

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 39

cccttctctt tgcgtgaatt ccatttc

27

&lt;210&gt; 40

&lt;211&gt; 22

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 40

gttgtgcccc taataaccag ag

22

&lt;210&gt; 41

&lt;211&gt; 3763

&lt;212&gt; DNA

&lt;213&gt; Oryza sp.

&lt;400&gt; 41

atggcggcga	ccatggcgtc	caacgcgcgcg	gctgcggcgcg	cggtgtccct	ggaccaggcc	60
gtggcggcgt	cggcggcgtt	ctcgctgcgcg	aagcagctgc	ggctgcccgc	cgcggcgcgc	120
gggggggatgc	gggtgcgggt	gcgggcgcckg	gggcggcggg	aggcggtggt	ggtggcgctcc	180
gcgtcgctcgt	cgtcggtggc	agcgcgcggcg	gcgaaggcgcg	aggagatcgt	gctccagccc	240
atcaggggaga	tctccggggc	ggttcagctg	ccagggtcca	agtcgctctc	caacaggatc	300
ctcctcctct	ccgccctctc	cgagggtgaga	cgcggatccc	ttcctcttgc	gtgaattcca	360
tttctggaga	tgagatttta	gggggtttat	taggtgaggt	ggctgtgttt	gtgaaatcct	420
aggaattatc	tctcaagtca	atctaacgat	gagatataac	tgaggttctg	gttttaatac	480
cacactcata	taaccaatth	attgaaacat	tttggtttgg	cataagaaac	tgettacgaa	540
ggatatgat	cctcctacat	gtcaggctac	ttaaattttca	cgacggtag	atccactcaa	600
aacaagtttc	ttaacgagtc	tggtgaggtc	tgttatgaaa	tttgtgtaaa	ctaaggcaac	660
tttgagggtt	tcgcactgta	ccaatgttat	gtttgaacat	tttgcaagca	gtgctttctc	720
ccaaaattat	gcaattttga	ggctcctcta	catcattata	attccccaat	acattgctct	780
ttattcttaa	tagctttgat	cgcgaaatth	aacattttta	ttcttgagct	gttattttgt	840
agcatcagtt	tatcatgagc	catgtttggt	actaaatata	caatcccttg	ggtttatctg	900
tttccaagca	tgctattaac	ttatcttaac	gtggacaaga	aactgatgcc	tgettacatt	960
gctattatth	caagcgggta	ttgatcctth	gacatgtgat	tgatcattth	ttttctctctg	1020
gttattaggg	cacaacagt	gtggacaact	tgctgaacag	tgaggatgth	cactacatgc	1080
ttgaggccct	gaaagccctc	gggtctctctg	tggaagcaga	taaagttgca	aaaagagctg	1140
tagtcgttgg	ctgtggtggc	aagtttctctg	ttgagaagga	tgcgaaagag	gaagtgcac	1200
tcttcttggg	gaacgctgga	actgcaatgc	gaccattgac	agcagccgtg	actgctgctg	1260
gtggaaatgc	aacgtatgth	ttttttttta	atgtttatga	aaatatgtat	ggaattcatg	1320
gggtatgtth	tatgacctth	ttctttacca	tcagttatgt	gcttgatgga	gtgccacgaa	1380
tgaggggagag	accgattggt	gacttggttg	tcgggttgaa	acaacttggt	gcggatgtcg	1440
actgtttcct	tggcactgaa	tgccccctg	ttcgtgtcaa	gggaattgga	ggacttctctg	1500
gtggcaaggt	tagttactcc	taaactgcat	cctttgtact	tctgtatgca	cctcaattct	1560

ttgtcaacct	tctgcattta	taaggaacat	tctatgatgc	aattcgacct	tacactgcac	1620
agtaacctga	aatgttttcat	gcttaaatcaa	tatgccatat	tcctgccaag	ctcaagcgag	1680
caataatttgt	ttgaatttgg	taccatattt	ttgtatattt	gggcattcct	ttttggtctt	1740
gatgtcttct	tttgaattag	catttaactg	aattacactc	aacagggttaa	gctctctggg	1800
tccatcagca	gtcagtaact	gagtgccttg	ctgatggctg	ctcctttggc	ccttggggat	1860
gtggagatcg	aaatcattga	caaactaatc	tccattcctt	acgttgaaat	gacattgaga	1920
ttgatggagc	gttttgggtg	gaaggcagag	cattctgata	gttgggacag	attctatatt	1980
aagggagggg	agaagtacaa	gtaagcttct	acctgcctta	ctgagctgaa	ttattcgggt	2040
gtttatgatt	aactccctaa	actaaccctt	tttcttttct	ttggcattga	cagatctcct	2100
ggaaatgcct	atgttgaagg	tgatgcctca	agcgcgagct	atttcttggc	tggtgctgca	2160
atcactggag	gcactgtgac	agttcaagg	tgtggtacga	ccagtttgca	ggtataactg	2220
tagtgcctgt	tttgacattc	taccgtttag	tcaagtttag	tcagtagtca	catattcaga	2280
atatagcaca	atctgtatta	tgccactggt	aatcaaatac	gcttgacctt	gagagtgcta	2340
tataccctag	cttaatcttc	aaactaaaca	gttctcttgt	ggcttgctgt	gctgttatgt	2400
tccttgacct	acatgttaat	attacagggt	gatgtcaaat	ttgctgaggt	acttgagatg	2460
atgggagcaa	aggttacatg	gactgacacc	agtgtaacgg	taactgggtcc	accacgtgag	2520
ccttatggga	agaaacacct	gaaagctgtt	gatgtcaaca	tgaacaaaat	gcctgatgtt	2580
gccatgaccc	ttgccgttgt	tgcactcttc	gctgatgggt	caactgctat	cagagatggg	2640
aaacattaa	gcctattata	cctgttctat	catactagca	attactgctt	agcattgtga	2700
aaaacaaat	aaccaaaact	tcttcaaaat	aacttagaaa	tataagaaa	gttcgttttg	2760
tgtggtaaac	agtactactg	tagtttcagc	tatgaagttt	gctgctggca	attttctgaa	2820
cggtttcagc	taaattgcat	gtttgttcat	catacttate	cattgtcttc	cacagtgggt	2880
tcctggagag	taaaggaaac	cgaaaggatg	gttgcaattc	ggaccgagct	aacaaaggta	2940
aattcattag	gtcccgtgtc	ctttcattct	tcaagtagtt	tgttcataag	ttgaattctc	3000
cttcaatgat	gtttaaattc	atcatcttct	tttttgggtg	tgtgccagct	gggagcatcg	3060
gttgaagaag	gtcctgacta	ctgcatcctc	acccacccgg	agaagctgaa	catcacggga	3120
atcgacacct	acgatgatca	caggatggcc	atggccttct	ccctcgctgc	ctgcgccgac	3180
gtgcccgtga	cgatcaggga	ccctggttgc	acccgcaaga	ccttccccc	ctacttcgac	3240
gttctaagca	ctttcgtcag	gaactgaact	gagcttttaa	aagagtgagg	tctaggttct	3300
gttgctctgtc	tgtccatcat	ccatgtgttg	actgttgagg	gaactcgttt	cttcttttct	3360
tcacgagatg	agtttttgtg	tgccgttaat	actagtttgt	agcaaaggct	gcgttacata	3420
aggtgatgag	aattgaggta	aatgagatc	tgtacactaa	attcattcag	actgttttgg	3480
cataaagaat	aatttggcct	tctgcgattt	cagaagctat	aaattgccat	ctcactaaat	3540
tctccttggt	cctcatggca	atgcaacgac	agtgtgaagc	actgaagccc	gtaatgctct	3600
atcaccacca	tgtacgacag	aacctatata	gtccatatgt	acaactcgag	tgttgtttga	3660
gtggccagca	aactggctga	ccaagccaca	cgagagagaa	tactataaac	tcaatcatat	3720
ataacaagcc	caagcaacat	tagacagaac	acaacaacac	tcg		3763

&lt;210&gt; 42

&lt;211&gt; 3763

&lt;212&gt; DNA

<213> *Oryza sp.*

&lt;400&gt; 42

atggcgggcga	ccatggcgctc	caacgccgcg	gctgcggcgg	cggtgtccct	ggaccaggcc	60
gtggcgggcgt	cgggcgcggt	ctcgctcgcg	aagcagctgc	ggctgcccgc	cgcggcgcgc	120
ggggggatgc	gggtgcgggt	gcgggcgckg	gggcggcggg	aggcggtggt	ggtggcgctcc	180
gcgtcgctcgt	cgtcggtggc	agcgccggcg	gcgaaggcgg	aggagatcgt	gctccagccc	240
atcaggggaga	tctccggggc	ggttcagctg	ccagggtcca	agtcgctctc	caacaggatc	300
ctcctcctct	ccgcctctct	cgaggtgaga	cgcggtatcc	ttcctcttgc	gtgaattcca	360
tttctggaga	tgagatttta	gggggtttat	taggtgaggt	ggctgtgttt	gtgaaatcct	420
aggaattatc	tctcaagtca	atctaacgat	gagatataac	tgaggttctg	gttttaatac	480
cacactcata	taaccaattt	attgaaacat	tttggtttgg	cataagaaac	tgcttacgaa	540
ggtatgatat	cctcctacat	gtcaggctac	taaattttca	cgacgggatg	atccactcaa	600
aacaagtttc	ttaacgagtc	tggtgaggtc	tgttatgaaa	tttgtgtaaa	ctaaggcaac	660
tttggagggt	tcgcactgta	ccaatgttat	gtttgaacat	tttgcaagca	gtgctttctc	720
ccaaaattat	gcaattttga	ggctcctcta	catcattata	attccccaat	acattgctct	780
ttattcttaa	tagctttgat	cgcgaaattt	aacattttta	ttcttgagct	gttattttgt	840
agcatcagtt	tatcatgagc	catgtttggg	actaaatata	caatcccttg	gggtttatttg	900
tttccaagca	tgtcattaac	ttatcttaat	gtggacaaga	aactgatgcc	tgcttacatt	960
gctattattt	caagcgggta	ttgatccttt	gacatgtgat	tgatcatttt	tttttctctg	1020

```

gttattagggt cacaacagtg gtggacaact tgctgaacag tgaggatggt cactacatgc 1080
ttgaggccctc gaaagccctc gggctctctg ttggaagcaga taaagttgca aaaagagctg 1140
tagtcgttgg ctgtggtggc aagtttcctg ttgagaagga tgcgaaagag gaagtgcac 1200
tcttcttggg gaacgctgga attgcaatgc gatcattgac agcagccgtg actgctgctg 1260
gtggaaatgc aacgtatgtt ttttttttta atgtttatga aaatatgtat ggaattcatg 1320
gggtatgttt tatgaccttt ttctttacca tcagttatgt gcttgatgga gtgccacgaa 1380
tgaggggagag accgattggt gacttggttg tggggttgaa acaacttggt gcggatgtcg 1440
actgtttcct tggcactgaa tgcccacctg tctgtgtcaa gggaattgga ggacttcctg 1500
gtggcaagggt tagttactcc taaactgcat cctttgtact tctgtatgca cctcaattct 1560
ttgtcaacct tctgcattta taaggaacat tctatgatgc aattcgacct tacactgcac 1620
agtaacttga aatgtttcat gcttaatcaa tatgccatat tcttgccaag ctcaagcgag 1680
caatatttgt ttgaatttgg taccatattt ttgtatatatt gggcattcct ttttggctct 1740
gatgtcttct tttgaattag catttaactg aattacactc aacaggttaa gctctctggt 1800
tccatcagca gtcagtactt gagtgccttg cctgtgctg ctcttttggc ccttggggat 1860
gtggagatcg aaatcattga caaactaate tccattcctt acgttgaaat gacattgaga 1920
ttgatggagc gttttggtgt gaaggcagag cattctgata gttgggacag attctatat 1980
aagggaaggc agaagtacaa gtaagcttct acctgcctta ctgagctgaa ttattcgggt 2040
gtttatgatt aactccctaa actaaccctt tttcttttct ttggcattga cagatctcct 2100
ggaaatgcct atgttgaagg tgatgcctca agcgcgagct atttcttggc tgggtgctgca 2160
atcactggag gcactgtgac agttcaagggt tgtggtacga ccagtttgca ggtataactg 2220
tagtgcttgt tttgacattc taccgtttag tcaagtttag tcagtagtca catattcaga 2280
atatagcaca atctgtatta tgccactgtt aatcaaatac gcttgaccta gagagtgcta 2340
tataccctag cttaatcttc aaactaaaca gttctcttgt ggcttgctgt gctgttatgt 2400
tccctgacct acatgttaat attacagggg gatgtcaaat ttgctgaggt acttgagatg 2460
atgggagcaa aggttacatg gactgacacc agtgtaaccc taactgggtc accacgtgag 2520
ccttatggga agaaacacct gaaagctgtt gatgtcaaca tgaacaaaat gcctgatgtt 2580
gccatgaccc ttgccgttgt tgcactcttc gctgatggtc caactgctat cagagatggg 2640
aaacattaag gcctattata cctgttctat catactagca attactgctt agcatttgtga 2700
caaaacaaat aaccaaactt tcttcaaaat aacttagaaa tataagaaaag gttcgttttg 2760
tgtggtaaac agtactactg tagtttcagc tatgaagttt gctgctggca attttctgaa 2820
cggtttcagc taaattgcat gtttgttcat catacttate cattgtcttc cacagtggct 2880
tccctggagag taaaggaaac cgaaaggatg gttgcaattc ggaccgagct aacaaaggta 2940
aattcattag gtcccgtgtc ctttcattct tcaagtagtt tgttcataag ttgaattctc 3000
cttcaatgat gtttaaatte atcatcttct tttttgggtg tgtgccagct gggagcatcg 3060
gttgaagaag gtccctgacta ctgcatcate accccaccgg agaagctgaa catcacggca 3120
atcgacacct acgatgatca caggatggcc atggccttct ccctcgctgc ctgcgcgac 3180
gtgcccggtg cgatcagggg ccttgggttg acccgcaaga ccttcccaa ctacttcgac 3240
gttctaagca ctttcgtcag gaactgaact gagcttttaa aagagtgagg tctaggttct 3300
gttgtctgtc tgtccatcat ccatgtgttg actgttgagg gaactcgttt cttcttttct 3360
tcacgagatg agtttttgtg tgccctgtaat actagtttgt agcaaaggct gcgttacata 3420
aggtgatgag aattgaggta aaatgagatc tgtacactaa attcattcag actggtttgg 3480
cataaagaat aatttggcct tctgcgattt cagaagctat aaattgccat ctcaactaa 3540
tctccttggg cctcatggca atgcaacgac agtgtgaagc actgaagccc gtaatgctct 3600
atcaccacca tgtacgacag aaccatatat gtccatatgt acaactcgag tgttgtttga 3660
gtggccagca aactggctga ccaagccaca cgagagagaa tactataaac tcaatcatac 3720
ataacaagcc caagcaacat tagacagaac acaacaacac tcg 3763

```

&lt;210&gt; 43

&lt;211&gt; 870

&lt;212&gt; DNA

&lt;213&gt; Zea mays

&lt;400&gt; 43

```

ttcagccttc gatgtggatg caacagcttc acaggattcc attaaatcgt agccattgtg 60
tcaaagtttg ctttgccaac gttatttatt tatttattta gaaaaccagc tttgaccagc 120
cgccctcttt acgtttggca caatttagct gaatccggcg gcatggcaag gtagactgca 180
gtgcagcgtg acccggtcgt gcccctctct agagataatg agcattgcat gtctaagtta 240
taaaaaatta ccacatattt ttttgtcaca cttgtttgaa gtgcagttta tctacttta 300
tacatatatt taaactttac tctacgaata tagtactaca ataatatcag 360
tgttttagag aatcatataa atgaacagtt agacatggct taaaggacaa ttgagtattt 420
tgacaacagg actctacagt tttatctttt tagtgtgcat gtgttctcct tttttttttg 480

```

```

caaatagctt  cacctatatata  atacttcatc  catttttatta  gtacatccat  ttaggggttta  540
gggttaatgg  tttttataga  ctaatttttt  tagtacatct  attttattct  attttagcct  600
ctaaatttaag  aaaactaaaa  ctctatttta  gtttttttat  ttaataattt  agatataaaa  660
tagaataaaa  taaagtgact  aaaaattaaa  caaataccct  ttaagaaatt  aaaaaaacta  720
aggaaacatt  tttcttggtt  cgagtagata  atgccagcct  gttaaacgcc  gtcgacgagt  780
ctaacggaca  ccaaccagcg  aaccagcagc  gtcgctcgg  gccaaagcgaa  gcagacggcg  840
cggcatctct  gtcgctgcct  ctggaccctt  870

```

<210> 44  
 <211> 1501  
 <212> DNA  
 <213> *Oryza* sp.

```

<400> 44
gatatccctc  agccgccttt  cactatcttt  tttgcccgag  tcattgtcat  gtgaaccttg  60
gcatgtataa  tcggtgaatt  gcgtcgattt  tcctcttata  ggtgggccaa  tgaatccgtg  120
tgatcgcgct  tgattggcta  gagatatgtt  tcttccttgt  tggatgtatt  ttcatacata  180
atcatatgca  tacaaatatt  tcattacact  ttatagaaat  ggtcagtaat  aaaccctatc  240
actatgtctg  gtgtttcatt  ttatttgctt  ttaaaccgaa  attgacttcc  tgattcaata  300
tttaaggatc  gtcaacgggtg  tgcagttact  aaattctggt  ttgtaggaac  tatagtaaac  360
tattcaagtc  ttcacttatt  gtgcactcac  ctctcgccac  atcaccacag  atgttattca  420
cgtctttaat  ttgaactaca  catcatattg  acacaatatt  ttttttaaat  aagcgattaa  480
aacctagcct  ctatgtcaac  aatgggtgtac  ataaccagcg  aagtttaggg  agtaaaaaac  540
atcgccctac  acaaagtctg  ctttaaaaaa  taaagagtaa  attttacttt  ggaccacctt  600
tcaaccaatg  tttcacttta  gaacgagtaa  ttttattatt  gtcactttgg  accacctca  660
aatctttttt  ccatctacat  ccaattttat  atgtcaaaga  aatggtctac  atacagctaa  720
ggagatttat  cgacgaatag  tagctagcat  actcgaggtc  attcatatgc  ttgagaagag  780
agtcgggata  gtccaaaata  aaacaaagggt  aagattacct  ggtcaaaagt  gaaaacatca  840
gttaaaagggt  ggtataaaagt  aaaatatcgg  taataaaagg  tggcccaaag  tgaaattttac  900
tcttttctac  tattataaaa  attgaggatg  tttttgtcgg  tactttgata  cgtcattttt  960
gtatgaattg  gtttttaagt  ttattcgctt  ttggaaatgc  atatctgtat  ttgagtcggg  1020
ttttaagttc  gtttgctttt  gtaaatacag  agggatttgt  ataagaaata  tctttaaaaa  1080
aaccatattg  ctaatttgac  ataatttttg  agaaaaatat  atattcaggc  gaattctcac  1140
aatgaacaat  aataagatta  aaatagcttt  ccccggttgc  agcgcatggg  tattttttct  1200
agtaaaaaa  aaagataaac  ttagactcaa  aacatttaca  aaaacaacct  ctaaagttcc  1260
taaagcccaa  agtgctatcc  acgatccata  gcaagcccaa  cccaacccaa  cccaacccaa  1320
cccacccag  tccagccaac  tggacaatag  tctccacacc  cccccactat  caccgtgagt  1380
tgtccgcag  caccgcacgt  ctgcgagcca  aaaaaaaaaa  aagaaagaaa  aaaaagaaaa  1440
agaaaaaaca  gcagggtgggt  ccgggtcgtg  ggggccggaa  acgcgaggag  gatcgcgagc  1500
a  1501

```

<210> 45  
 <211> 982  
 <212> DNA  
 <213> *Oryza* sp.

```

<400> 45
gttggttggt  gagagtgaga  caccgacgga  acggaaggag  aaccacgccc  cttggatttt  60
tcttttttac  cttttcaaat  ttttaattta  aaaataaaac  catttttaaa  acttatcttc  120
aaatacaaat  ctttttaaaa  cactaacacg  tgacacacag  cgggcacgtc  acccaaacgg  180
gcgtgacaat  attgttttgc  cacaccaatc  cagctgggtg  ggacaaaatg  ttcataatatt  240
gaaaaataaaa  tttaaaacaa  tttatatatt  ttatctatat  cattataaaa  attgaagatg  300
tttttaccgg  tattttgtta  ctattttgtg  catgagtcgg  tttttaagtt  tgttcgcttt  360
tggaataaca  tatccgtatt  tgagtatgtt  ttttaagttc  ttcgtttttt  gaaatacaaa  420
aggaatcgta  aaataaatct  attttaaaaa  actcgcatgc  taacttgaga  cgatcgaact  480
gctaattgca  gtcataatt  ttccaaaaaa  aaatatatcc  aaacgagttc  ttatagtaga  540
tttcacctta  attaaaacat  ataaatgttc  acccggtaca  acgcacgagt  atttttataa  600
gtaaaattaa  aagtttaaaa  taaataaaaa  tcccgcacc  acggcgcgat  ggtaaaaggg  660
ggacgcttct  aaacggggcc  ggcacgggac  gatcgggccc  gaacccggcc  catctaaccg  720
ctgtagggcc  accgcccacc  aatccaactc  cgtactacgt  gaagcgctgg  atccgcaacc  780
cgtaagcag  tccacacgac  tcgactcgac  tcgcgcactc  gccgtggtag  gtggcaaccc  840

```

```

ttcttctctcc tctattttctt cttcttctctc ccttctctcgc ctcaccacac caaccgcacc 900
aaccccaacc ccgcgcgcgc tctccctct cccctcccac caaccacc ccatcctccc 960
gacctccacg ccgcgcggcaa tg 982

```

<210> 46  
 <211> 435  
 <212> DNA  
 <213> *Oryza* sp.

```

<400> 46
ttaattaaaa catataaatg ttcacccggg acaacgcacg agtatttttta taagtaaaat 60
taaaagtttta aaataaataa aaatcccgcg accacggcgc gatggtaaaa gggggacgct 120
tctaaacggg ccgggcacgg gacgatcggc cccgaacccg gcccatctaa ccgctgtagg 180
cccaccgccc accaatccaa ctccgtacta cgtgaagcgc tggatccgca acccgtaaag 240
cagtcacac gactcgactc gactcgcgca ctgcgcgtgg taggtggcaa cccttcttcc 300
tcctctatctt cttcttcttc ctcccttctc cgctcacca caccaaccgc accaacccca 360
acccgcgcgc cgctctcccc tctccctcc caccaacccc accccatcct cccgacctcc 420
acgcgcgcgg caatg 435

```

<210> 47  
 <211> 1343  
 <212> DNA  
 <213> *Oryza* sp.

```

<400> 47
gccacaccaa tccagctggg gtggacaaaa tgttcatata ttgaaaataa aattttaaac 60
aatttatatt ttttatctat atcattataa aaattgaaga tgtttttacc ggtattttgt 120
tactcatttg tgcagtagtc ggttttttaag tttgttcgct tttggaaata catatccgta 180
tttgagtagt tttttaagtt cgttcgtttt ttgaaataca aaaggaatcg taaaataaat 240
ctatttttaa aaactcgcat gctaacttga gacgatcgaa ctgctaattg cagctcataa 300
ttttccaaaa aaaaatatat ccaaacgagt tcttatagta gatttcacct taattaaaac 360
atataaatgt tcacccggta caacgcacga gtatttttat aagtaaaatt aaaagtttta 420
aataaataaa aatcccgcga ccacggcgcg atggtaaaag ggggacgctt ctaaaccgggc 480
cgggcacggg acgatcggcc ccgaacccgg cccatctaac cgctgtaggc ccaccgcca 540
ccaatccaac tccgtactac gtgaagcgct ggatccgcaa cccgttaagc agtccacacg 600
actcgactcg actcgcgcac tcgcgcgtgg aggtggcaac ccttcttctc cctctatttc 660
ttcttcttcc tcccttctcc gcctcaccac accaaccgca ccaaccccaa cccgcgcgcg 720
gctctccctc ctcccttccc accaacccca ccccatcctc ccgacctcca cgcgcgcggc 780
aggatcaagt gcaaagggtc gccttggttc tectctgtct ctgatctga ctaatcttgg 840
tttatgattc gttgagtaat tttggggaaa gctagcttcg tccacagttt ttttttcgat 900
gaacagtgcc gcagtggcgc tgatcttgta tgctatcctg caatcgtggg gaacttattt 960
cttttatacc cttcactccc atgaaaaggt tagtaatctt tctcgatgta acatcgcca 1020
gcactgctat taccgtgtgg tccatccgac agtctggctg aacacatcat acgatattga 1080
gcaaagatct atcctccctg ttctttaatg aaagacgtca ttttcatcag tatgatctaa 1140
gaatggtgca acttgcaagg aggcgtttct ttctttgaat ttaactaact cgttgagtagg 1200
ccctgtttct cggacgtaag gcctttgctg ctccacacat gtccattcga attttaccgt 1260
gtttagcaag agcgaaaagt ttgcactctg atgatttagc ttgactatgc gattgctttc 1320
ctggaccgct gcagctgcgg atg 1343

```

<210> 48  
 <211> 538  
 <212> DNA  
 <213> *Zea mays*

```

<400> 48
gtccgccttg tttctcctct gtctcttgat ctgactaatc ttggtttatg attcgttgag 60
taattttggg gaaagctagc ttcgctccaca gttttttttt cgatgaacag tgccgcagtg 120
gcgctgatct tgtatgctat cctgcaatcg tggggaactt atttctttta tacccttcac 180
tcccataaaa aggctagtaa tctttctcga tgtaacatcg tccagcactg ctattaccgt 240
gtggtccatc cgacagtctg gctgaacaca tcatacgata ttgagcaaag atctatcctc 300
cctgttcttt aatgaaagac gtcattttca tcagtatgat ctaagaatgt tgcaacttgc 360

```

```
aaggaggcgt ttctttcttt gaatttaact aactcgttga gtggccctgt ttctcggacg 420
taaggccttt gctgctccac acatgtccat tcgaatttta ccgtgttttag caagagcgaa 480
aagtttgcac cttgatgatt tagcttgact atgcgattgc ttcctggac ccgtgcag 538
```

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01572

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 C12N9/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BIOSIS

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GASSER C S ET AL: "STRUCTURE EXPRESSION AND EVOLUTION OF THE 5 ENOLPYRUVYLSHIKIMATE-3-PH OSPHATE SYNTHASE GENES OF PETUNIA AND TOMATO" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 263, no. 9, 1988, pages 4280-4287, XP002145760 ISSN: 0021-9258 see esp. p.4283, fig.3 ---	4, 19, 32
X	WO 98 44140 A (MCELROY DAVID; STEPHENS MICHAEL A; SPENCER T MICHAEL ; MUMM RITA (U) 8 October 1998 (1998-10-08) the whole document --- -/--	4, 19, 32

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

25 August 2000

Date of mailing of the international search report

12/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Kania, T



## INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01572

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 91 04323 A (MONSANTO CO) 4 April 1991 (1991-04-04)  see the whole document; esp. p.9 l.30 - p.10 l.12; p.33 l.10-26 ---	4, 19, 32, 33, 35, 37, 39, 41-44, 46-50
X	WO 92 06201 A (MONSANTO CO) 16 April 1992 (1992-04-16)  see the whole document; esp. p.9 l.26 - p.10 l.10; p.32 l.20 - p.33 l.4; p.65 l.12-19; examples ---	4, 19, 32, 33, 35, 37-39, 41-44, 46-52 20-22
A	EP 0 459 643 A (COMMW SCIENT IND RES ORG ;LUBRIZOL GENETICS INC (US)) 4 December 1991 (1991-12-04) the whole document ---	5-23
A	WO 97 20056 A (CAMBRIDGE ADVANCED TECH ;GRAY JOHN CLINTON (GB); SANDHU JAGDEEP SI) 5 June 1997 (1997-06-05) the whole document ---	5-23
A	WO 97 04103 A (LEBRUN MICHEL ;SAILLAND ALAIN (FR); FREYSSINET GEORGES (FR); RHONE) 6 February 1997 (1997-02-06) see the whole document; esp. p.2 l.7-16; examples ---	1-52
A	GB 2 326 163 A (ZENECA LTD) 16 December 1998 (1998-12-16) see esp. p.6 l.11 - p.11 l.11 ---	1-52
A	WO 99 10513 A (DANIEL HENRY ;UNIV AUBURN (US)) 4 March 1999 (1999-03-04) the whole document ---	1-52
A	WO 98 39419 A (ASGROW SEED CO) 11 September 1998 (1998-09-11) see esp. example 1 ---	25, 26, 34
A	KISHIMA Y AND SANO Y. : "Oryza sativa mRNA for EPSP-synthase; AC AB016765" EBI DATABASE, 12 August 1998 (1998-08-12), XP002145761 the whole document -----	1-3

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 00/01572

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9844140 A	08-10-1998	US 6040497 A AU 6882298 A EP 0975778 A	21-03-2000 22-10-1998 02-02-2000
WO 9104323 A	04-04-1991	US 5310667 A AU 640179 B AU 6638190 A BR 9007550 A CA 2059266 A EP 0409815 A EP 0483287 A NZ 234532 A ZA 9005570 A	10-05-1994 19-08-1993 18-04-1991 23-06-1992 18-01-1991 23-01-1991 06-05-1992 29-01-1992 29-05-1991
WO 9206201 A	16-04-1992	US 5866775 A AT 118820 T AU 8719291 A CA 2090617 A DE 69107621 D EP 0550633 A JP 6501615 T	02-02-1999 15-03-1995 28-04-1992 29-03-1992 30-03-1995 14-07-1993 24-02-1994
EP 0459643 A	04-12-1991	AU 643521 B AU 7710791 A CA 2042831 A CN 1063506 A JP 7067645 A US 5290924 A	18-11-1993 21-11-1991 19-11-1991 12-08-1992 14-03-1995 01-03-1994
WO 9720056 A	05-06-1997	AU 7635696 A BR 9611657 A CA 2236166 A EP 0870043 A NZ 322576 A	19-06-1997 23-02-1999 05-06-1997 14-10-1998 28-10-1999
WO 9704103 A	06-02-1997	FR 2736926 A AU 6619196 A BG 102247 A BR 9609792 A CA 2223875 A CN 1196088 A CZ 9800174 A EP 0837944 A HU 9900463 A JP 11510043 T PL 324572 A SK 6498 A	24-01-1997 18-02-1997 30-04-1999 16-03-1999 06-02-1997 14-10-1998 17-06-1998 29-04-1998 28-05-1999 07-09-1999 08-06-1998 04-11-1998
GB 2326163 A	16-12-1998	AU 7541498 A EP 1017825 A WO 9854330 A	30-12-1998 12-07-2000 03-12-1998
WO 9910513 A	04-03-1999	AU 8457398 A EP 1002115 A	16-03-1999 24-05-2000
WO 9839419 A	11-09-1998	AU 6758198 A	22-09-1998